

Time-optimal navigation through quantum wind

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 New J. Phys. 17 033048

(<http://iopscience.iop.org/1367-2630/17/3/033048>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.83.1.242

This content was downloaded on 13/04/2015 at 08:51

Please note that [terms and conditions apply](#).



PAPER

Time-optimal navigation through quantum wind

OPEN ACCESS

RECEIVED

25 November 2014

REVISED

2 February 2015

ACCEPTED FOR PUBLICATION

18 February 2015

PUBLISHED

31 March 2015

Content from this work
may be used under the
terms of the [Creative
Commons Attribution 3.0
licence](#).

Any further distribution of
this work must maintain
attribution to the
author(s) and the title of
the work, journal citation
and DOI.

Dorje C Brody^{1,2}, Gary W Gibbons³ and David M Meier¹¹ Department of Mathematics, Brunel University London, Uxbridge UB8 3PH, UK² St Petersburg National Research University of Information Technologies, Mechanics and Optics, 49 Kronverksky Avenue, St Petersburg 197101, Russia³ Department of Applied Mathematics and Theoretical Physics, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WA, UKE-mail: Dorje.Brody@brunel.ac.uk

Keywords: controlled quantum dynamics, Zermelo navigation, Finsler geometry


Abstract

The quantum navigation problem of finding the time-optimal control Hamiltonian that transports a given initial state to a target state through quantum wind, that is, under the influence of external fields or potentials, is analyzed. By lifting the problem from the state space to the space of unitary gates realizing the required task, we are able to deduce the form of the solution to the problem by deriving a universal quantum speed limit. The expression thus obtained indicates that further simplifications of this apparently difficult problem are possible if we switch to the interaction picture of quantum mechanics. A complete solution to the navigation problem for an arbitrary quantum system is then obtained, and the behaviour of the solution is illustrated in the case of a two-level system.

With the advances in the implementation of quantum technologies, the theoretical understanding of controlled quantum dynamics and, in particular, of their limits, is becoming increasingly important. One aspect of such limits that has been investigated extensively in the literature concerns the time-optimal manoeuvring of quantum states [1–18]. If the time-evolution is unconstrained (apart from a bound on the energy resource), then this amounts to finding the time-independent Hamiltonian that generates maximum speed of evolution. However, in general there can be a range of constraints that prohibits the implementation of such an elementary protocol, and various optimizations will have to be applied to determine time-dependent Hamiltonians that generate the dynamics achieving required tasks.

An important class of problems arising in this context is the identification of the time-optimal quantum evolution under the influence of external fields or potentials that cannot be easily eliminated in a laboratory. Solutions to such problems are indeed relevant to practical implementations of time-optimal controlled quantum dynamics because in real laboratories external influences (e.g. electromagnetic fields) are typically present. Problems of this kind can be thought of as representing the quantum counterpart of the classical Zermelo navigation problem of finding the time-optimal control that takes a ship from one location to another, under the influence of external wind or currents [19, 20]. Within the context of quantum Zermelo problems, there are two distinct questions that arise, namely, finding the time-optimal Hamiltonian (i) that generates a required unitary gate; and (ii) that transports a given initial state to a required target state. In the context of constructing a unitary gate, this problem was formulated in [21], and solved more recently in [22, 23]. The construction of the time-optimal Hamiltonian that transports a given initial state to a target state, on the other hand, is *a priori* more challenging (see also [24]), because (as shown below) this involves a variational problem with free boundaries. The purpose of the present paper is to derive the full solution to this latter problem for an arbitrary quantum system.

One way of addressing the quantum-state navigation problem is to work in the space of rays through the origin of the underlying Hilbert space (i.e. the complex projective space, or simply the ‘state space’). Then the Schrödinger evolution generated by the background field gives rise to a Hamiltonian vector field on the state space, and the task at hand can be formulated as a Hamiltonian control problem, the solution of which is typically difficult to obtain. We are nevertheless able to go forward by exploiting certain subtle geometric



structures of the unitary group, which in turn allows us to efficiently make use of the results obtained in [22, 23]. A remarkable feature of the solution that we obtain is that it is strikingly reminiscent of the analysis of quantum dynamics in the interaction-picture of Schwinger and Tomonaga [25]. On account of this observation we are able to revisit our analysis by employing a physical symmetry argument, rather than purely variational reasoning. As a consequence, a considerably simpler derivation of the solution emerges. In fact, it turns out that by invoking the line of thinking behind the Schwinger–Tomonaga theory, a straightforward solution can be found of closely-related and apparently difficult problems in geometry [26, 27]. For an illustration we conclude the paper by presenting examples in the case of a two-level system.


Let us begin by stating more explicitly the Zermelo navigation problem for quantum states: For a given initial state $|\psi_I\rangle$ and a final target state $|\psi_F\rangle$, the task of an experimentalist is to find the control Hamiltonian $\hat{H}_1(t)$ such that the total Hamiltonian $\hat{H}(t) = \hat{H}_0 + \hat{H}_1(t)$ will generate the transformation $|\psi_I\rangle \rightarrow |\psi_F\rangle$ in the shortest possible time. Here \hat{H}_0 represents the Hamiltonian of the background field or potential that cannot be manipulated. Since the objective is to realize a time-optimal transformation (navigation), it is assumed that the available energy resource, which evidently has to be bounded, will be consumed fully. This translates into the ‘full throttle’ condition that the squared magnitude of the evolution speed generated by the control Hamiltonian \hat{H}_1 , which is given on account of the Anandan–Aharonov relation [28] by four times the variance $4\Delta H_1^2$ of \hat{H}_1 , is held fixed at all times at the maximum attainable value.

To illustrate the nature of the task involved, let us first consider a more elementary problem of Zermelo navigation on the plane \mathbb{R}^2 , with constant (in space and in time) wind, represented by the vector \vec{w} . Suppose that, departing from the origin o the desired destination is given by the endpoint of a vector $\vec{\xi}$. To work out the time it takes to reach the destination, consider a circle whose radius is determined by the distance that can be reached with full speed over one unit of time. In the absence of wind this unit circle is clearly centred at the origin. However, under the influence of the wind the centre of the circle is shifted by \vec{w} . For the moment let us assume that the wind is not dominant, i.e. $|\vec{w}| < 1$ so that the origin remains inside the shifted circle. This configuration is schematically illustrated in figure 1. Under the influence of the wind, therefore, to reach the endpoint ξ of the vector $\vec{\xi}$, it suffices to determine the distance $|\vec{o\rho_\xi}|$, where ρ_ξ is the point of intersection of the vector $\vec{o\xi}$ with the shifted circle. This follows on account of the fact that since it takes one unit of time to reach ρ_ξ , it will take $F(\xi) = |\vec{o\xi}|/|\vec{o\rho_\xi}|$ units of time to reach ξ . Elementary algebra then shows that the required time $F(\xi)$ is given by the expression

$$F(\xi) = \frac{\sqrt{\langle \vec{w}, \vec{\xi} \rangle^2 + |\vec{\xi}|^2 (1 - |\vec{w}|^2)} - \langle \vec{w}, \vec{\xi} \rangle}{1 - |\vec{w}|^2}, \quad (1)$$

where $\langle \vec{w}, \vec{\xi} \rangle = \vec{w} \cdot \vec{\xi}$ is the Euclidean inner product, $|\vec{w}|^2 = \langle \vec{w}, \vec{w} \rangle$, and $|\vec{\xi}|^2 = \langle \vec{\xi}, \vec{\xi} \rangle$.

The foregoing analysis on \mathbb{R}^2 extends naturally to the general case of navigation on a Riemannian manifold of arbitrary dimension, albeit the Euclidean inner product $\langle \cdot, \cdot \rangle$ has to be replaced by the Riemannian one. This is because the total journey time can be decomposed into a large number of infinitesimal journey times, each on a



tangent space of the manifold. In the case of optimal quantum navigation, we recall that the space of pure states is equipped with the unitary-invariant Fubini–Study metric [29–31]. The wind \vec{w} is then replaced by the infinitesimal Hamiltonian symplectic flow generated by \hat{H}_0 , whereas $\vec{\xi}$ is some vector tangent to quantum state space. The total journey time T along a path on state space is given by the integral of $F(\xi)$, where $\vec{\xi}$ is the velocity vector. In particular, for the Zermelo problem, $\vec{\xi}$ is generated by $\hat{H}(t) = \hat{H}_0 + \hat{H}_1(t)$, and this implies that $F(\xi)$ in the quantum context is given by a function of the variance ΔH_0^2 of \hat{H}_0 , the variance ΔH_1^2 of \hat{H}_1 , and the covariance $\langle \hat{H}_0 \hat{H}_1 \rangle$ of the two Hamiltonians. The optimal control is obtained by minimizing T over all $\hat{H}_1(t)$, subject to the full throttle constraint $4\Delta H_1^2 = 1$. The no-dominance condition on the wind is required in the case of a navigation problem on an open manifold such as \mathbb{R}^n , since otherwise the target may never be reached. In the quantum context we are able to relax the requirement $4\Delta H_0^2 < 1$, since the manifold of pure states is compact, which means that if the wind against the shortest path is too strong, one can always go ‘the other way around’ to reach the target state (note incidentally that $F(\xi)$ remains well defined in the limit $|\vec{w}| \rightarrow 1$ since $F(\xi) \rightarrow |\vec{\xi}|^2/2\langle \vec{w}, \vec{\xi} \rangle$).

The Hamiltonian control problem specified above is not straightforward to solve directly, and this leads us to employ an alternative approach based on the idea of lifting the problem from state space to the unitary group acting on the states. In fact, from a physical point of view this is more appealing, since, although we are seeking the time-optimal path on state space, physical implementations are always carried out by constructing the Hamiltonian that generates the desired unitary operator.

With this in mind, we proceed by considering a fibre space above the state space with the property that for any state $|\psi\rangle$, the fibre above it consists of elements $\{\hat{u}\}$ of the special unitary group fulfilling the condition that $\hat{u}|\psi_I\rangle = |\psi\rangle$. Although our solution is applicable to both finite and infinite dimensional Hilbert spaces, to simplify the discussion let us assume that the Hilbert space is of finite dimension $n+1$. The totality of such fibres, when bundled together, then forms the group $\text{SU}(n+1)$, which acts on the state space (the complex projective space $\mathbb{C}\mathbb{P}^n$). This configuration is illustrated in figure 2. Let \hat{u}_F be an element of the fibre above the target state $|\psi_F\rangle$, i.e. $\hat{u}_F|\psi_I\rangle = |\psi_F\rangle$. Then the problem of finding the time-optimal transformation $|\psi_I\rangle \rightarrow |\psi_F\rangle$ translates into the problem of constructing the unitary gate \hat{u}_F from the identity element $\hat{e} \in \text{SU}(n+1)$ in the shortest possible time, under the influence of an external field. This translation of the problem helps us because the optimal Zermelo navigation for a specified unitary gate has been worked out recently [22, 23], and we can make use of the result in deducing the optimal control Hamiltonian for the state transfer; however, we encounter two difficulties: (a) the ‘target’ unitary gate \hat{u}_F is not unique since *a priori* it can be any element of the fibre above $|\psi_F\rangle$. In other words, if \hat{v}_F is any unitary operator that leaves the state $|\psi_F\rangle$ invariant, then the unitary gate $\hat{v}_F\hat{u}_F$ also satisfies the required condition $\hat{v}_F\hat{u}_F|\psi_I\rangle = |\psi_F\rangle$ so we have to deal with a free final boundary; (b) the ‘full throttle’ condition $4\Delta H_1^2 = 1$ on state space is concerned with the evolution speed, whereas the condition used in [21–23] for unitary gates appears to be distinct since it concerns the Hilbert–Schmidt norm $2\text{tr}(\hat{H}_1^2) = 1$. In

what follows we shall show that these two apparent issues evaporate once the relevant ‘horizontality condition’ is imposed on the control Hamiltonian.

To see this we begin by noting that since $\hat{e}|\psi_I\rangle = |\psi_I\rangle$, the identity element \hat{e} belongs to the fibre above the initial state $|\psi_I\rangle$. In fact, the fibre above $|\psi_I\rangle$ coincides with the totality of unitary group elements $\{\hat{v}_I\}$ that leave the initial state $|\psi_I\rangle$ invariant. Intuitively, to reach from \hat{e} a target gate \hat{u}_F above $|\psi_F\rangle$ in a timely manner we would like to move away from \hat{e} as quickly as possible, and this is achieved by manoeuvring *horizontally*, i.e. towards a direction that is orthogonal to the fibre. That is to say, the choice of the initial control Hamiltonian $\hat{H}_1(0)$ has to be such that $\text{tr}(\hat{H}_1(0)\hat{H}_I) = 0$ for all Hamiltonians \hat{H}_I that leave the initial state invariant (i.e. $\hat{H}_1(0)$ has to be orthogonal to all generators of \hat{v}_I). This suggests that the horizontality condition $\text{tr}(\hat{H}_1(0)\hat{H}_I) = 0$ gives rise to the maximum evolution speed. To show this more explicitly we shall derive a kind of universal quantum speed limit. The statement of the result we shall establish is as follows: The squared speed of the evolution of a quantum state generated by a Hamiltonian \hat{H} , as defined by Anandan and Aharonov [28], is bounded above by twice the Hilbert–Schmidt norm $\text{tr}(\hat{H}^2)$ of the Hamiltonian, and the bound is attained if \hat{H} is horizontal. The implication of this result is that under the horizontality requirement the norm condition and the maximum speed condition are equivalent, and this resolves the issue (b) raised above. As a consequence, we are able to borrow the results of [22, 23] to deduce that the time-optimal control Hamiltonian $\hat{H}_1(t)$ is necessarily of the form

$$\hat{H}_1(t) = e^{-i\hat{H}_0 t} \hat{H}_1(0) e^{i\hat{H}_0 t}. \quad (2)$$

Notice that the horizontality condition is preserved under the adjoint action (2), which is indeed required since the full throttle condition has to be maintained throughout the operation.

The horizontality condition alone, of course, does not fix $\hat{H}_1(0)$ uniquely, since the constraint $\text{tr}(\hat{H}_1(0)\hat{H}_I) = 0$ on $\hat{H}_1(0)$ leaves $2n$ degrees of freedom. However, by demanding that the required gate should generate the target state $|\psi_F\rangle$, which imposes $2n$ conditions, we are able to select a unique initial control Hamiltonian $\hat{H}_1(0)$. In other words, the freedom in the boundary is eliminated by the horizontality condition $\text{tr}(\hat{H}_1(0)\hat{H}_I) = 0$ along with the boundary condition $\hat{u}_F|\psi_I\rangle = |\psi_F\rangle$, and this resolves the issue (a) raised above.

Before we turn to determining the initial control Hamiltonian $\hat{H}_1(0)$, and the amount of time T that is needed to reach the target state under the optimal control, let us discuss the claim above on the universal speed limit (see also [32]). From a physical point of view the claim is plausible if we make note of the following observation. Suppose that the Hamiltonian contains a vertical component tangent to the fibre that leaves the state below invariant. In this case, although energy is scarce, the experimentalist will be consuming energy that produces no work (except for shifting the overall phase). This is clearly not energy efficient. It follows that the optimal performance is ensured by eliminating vertical components altogether. A more precise derivation, which essentially follows from a standard result on complex projective spaces outlined, e.g., in Kobayashi and Nomizu [33] (see section 11.10), is given in the appendix.


Having established the ‘geodesic’ curve (2) that minimizes the action $T = \int F ds$ we are now in the position to identify the initial control $\hat{H}_1(0)$ so that the target state is reached. To this end we observe that the form (2) of the solution is indicative of the analysis in the interaction picture of quantum mechanics, if we view \hat{H}_0 as the ‘free’ and $\hat{H}_1(0)$ as the ‘interaction’ Hamiltonian. Alternatively stated, if we change from the ‘rest frame’ to the ‘moving frame’ on the state space according to the wind \hat{H}_0 , then without the control Hamiltonian the initial state remains fixed, while the target state moves against the wind according to $|\psi_F(t)\rangle = e^{i\hat{H}_0 t} |\psi_F\rangle$. Evidently, the state $|\psi_F(t)\rangle$ thus defined is the solution to the (nonrelativistic version of) the Tomonaga–Schwinger equation. In particular, standard results in the interaction-picture analysis [25] show that the time-ordered product of the unitary group generated by the Hamiltonian $\hat{H}(t) = \hat{H}_0 + e^{-i\hat{H}_0 t} \hat{H}_1(0) e^{i\hat{H}_0 t}$ gives rise to the evolution equation of the form

$$|\psi(t)\rangle = e^{-i\hat{H}_0 t} e^{-i\hat{H}_1(0)t} |\psi_I\rangle. \quad (3)$$

This result can also be verified directly (see also [22]) by differentiation: writing $\hat{u}(t) = e^{-i\hat{H}_0 t} e^{-i\hat{H}_1(0)t}$, we find $\partial_t \hat{u}(t) = -i\hat{H}(t) \hat{u}(t)$. As indicated above, to fix the initial condition $\hat{H}_1(0)$ we need to make use of the boundary condition, which implies that

$$e^{-i\hat{H}_0 T} e^{-i\hat{H}_1(0)T} |\psi_I\rangle = |\psi_F\rangle \quad (4)$$

must hold for some T that is to be determined. Observe that there are $2n$ unknowns in $\hat{H}_1(0)$, and we have an additional unknown parameter T . The boundary condition (4) gives rise to $2n$ constraints. Together with the norm condition $2\text{tr}(\hat{H}_1^2) = 1$ we are thus able to fix all the unknowns, and this in turn solves the quantum-state Zermelo navigation problem.



To proceed with this let us rewrite (4) in the form

$$e^{-i\hat{H}_1(0)T} |\psi_I\rangle = e^{i\hat{H}_0T} |\psi_F\rangle. \quad (5)$$

Although the transformation from (4) to (5) is in itself trivial, it sheds a different light on the problem at hand. As indicated above, if we transform to the moving frame generated by the wind such that the initial state remains stationary, then the objective becomes ‘hitting’ a moving target $e^{i\hat{H}_0t} |\psi_F\rangle$ as quickly as possible. Exactly how long it will take to achieve this task depends on how fast the motion towards the target can be, but for a fixed speed the shortest time T achievable is attained by following the geodesic path on the state space joining $|\psi_I\rangle$ and $e^{i\hat{H}_0T} |\psi_F\rangle$. On the other hand, the curve $|\psi(t)\rangle = e^{-i\hat{H}_1(0)t} |\psi_I\rangle$ joining $|\psi_I\rangle$ and $e^{-i\hat{H}_1(0)T} |\psi_F\rangle$ is a geodesic on the Fubini–Study manifold if and only if $\hat{H}_1(0)$ is horizontal, and this is indeed the situation we have in (5). To determine $\hat{H}_1(0)$ explicitly, let $\theta = \theta(T)$ be the angular distance between $|\psi_I\rangle$ and $e^{i\hat{H}_0T} |\psi_F\rangle$ defined by the relation:

$$\left| \langle \psi_I | e^{i\hat{H}_0T} |\psi_F\rangle \right| = \cos \frac{1}{2}\theta. \quad (6)$$

Evidently, the angular distance $\theta(T) = \omega T$ is given by the speed ω of the evolution generated by $\hat{H}_1(0)$ multiplied by the duration T of the journey, but the evolution speed is fixed on account of the condition $2\text{tr}(\hat{H}_1^2) = 1$ to $\omega = 1$, so we deduce that


$$2 \arccos \left| \langle \psi_I | e^{i\hat{H}_0T} |\psi_F\rangle \right| = T. \quad (7)$$

Since \hat{H}_0 , $|\psi_I\rangle$, and $|\psi_F\rangle$ are given, we see therefore that the smallest positive root of (7) uniquely determines T . Once T is fixed, the optimal control Hamiltonian at time zero can be obtained explicitly by making use of the result in [10]:

$$\hat{H}_1(0) = \frac{i \left(|\psi_I\rangle \langle \psi_F| e^{-i\hat{H}_0T} - e^{i\hat{H}_0T} |\psi_F\rangle \langle \psi_I| \right)}{2 \sin\left(\frac{1}{2}T\right)}. \quad (8)$$

Note that without loss of generality we can adjust the overall phase such that $\langle \psi_I | e^{i\hat{H}_0T} |\psi_F\rangle$ is real. With this convention, which is assumed in (8), one can easily verify that this Hamiltonian is indeed horizontal.

This completes the derivation of the solution to the quantum-state Zermelo navigation problem: the optimal control Hamiltonian is given by (2), where $\hat{H}_1(0)$ is given by (8) and where the parameter T is determined by (7). To gain further intuition on the behaviour of the solution, in figure 3 we plot the trajectories of the quantum state generated by the Hamiltonian $\hat{H}(t) = \hat{H}_0 + \hat{H}_1(t)$ in the case of a two-level system. In particular, we show the time-optimal paths for a range of wind strengths, indicating for instance that if the head wind is too strong, then the optimal strategy is to switch the direction of the manoeuvre to turn the headwind into a tailwind. To investigate the minimum journey times associated with different wind strengths we have also considered a one-



parameter family of winds $\epsilon\hat{H}_0$, corresponding to the setup shown in figure 3, and determined the optimal time $T(\epsilon)$ for a range of ϵ ; the result is shown in figure 4.

In summary, we have been able to deduce in closed form the solution to the quantum Zermelo problem in complete generality. Since the result obtained takes a rather simple form, we expect that a practical implementation is feasible in a number of realistic situations. Of course, for some applications the controllable degrees of freedom in the Hamiltonian can be limited, in which case additional constraints will have to be imposed in finding the optimal control—some steps towards this direction in the context of unitary gates have been initiated in [22]. In fact, further progress can again be made by switching to the interaction picture: If we write $\{f_k(\hat{H}_1) = 0\}_{k=1,\dots,N}$ for the N constraints on the control Hamiltonian, then in the moving frame the constraints will take for each k the form $f_k(e^{-i\hat{H}_0 t}\hat{h}_k e^{i\hat{H}_0 t}) = 0$, where $\hat{h}_k = e^{i\hat{H}_0 t}\hat{H}_1 e^{-i\hat{H}_0 t}$ is the generator of the dynamics $i\partial_t|\psi\rangle = \hat{h}|\psi\rangle$. The analysis of [8], for instance, can then be applied with essentially just one modification, namely, that the target state is moving. The initial control Hamiltonian can then be identified by adapting the strategy proposed in [18].

It is worth noting that the use of the interaction picture, or equivalently the switch to the moving frame, which allowed us to simplify the problem considerably, has implications beyond the quantum navigation problem considered here. In this connection we remark that if we endow the quantum state space with the metric of the form (1), then the resulting space is known as a Randers space (see [34]), which is an example of a Finsler space (for $|w| \rightarrow 1$ the space reduces to a Kropina space). In the literature of Finsler geometry it is known that every Zermelo navigation problem on a Riemannian manifold can be solved by finding geodesics of the corresponding Randers space [26]. The classification of spaces in terms of their curvatures is of particular importance in geometry, and in the case of Randers spaces the identification of constant curvature metrics has been obtained by solving the associated navigation problem when the wind is a conformal vector field with constant scale factor [27] (the field generated by a unitary motion belongs to this class, with vanishing scale factor). However, and with hindsight, the main theorem of [27] can be proven in essentially two lines by adopting the interaction picture, and this shows that in the present context a physical intuition can offer insights into a purely mathematical question. We conclude by remarking that an interesting generalization of the present problem is to consider the case under which the external influences contain noise. In such a situation, the idea of reaching a target pure state is no longer tenable, whereas reaching a particular mixed state might be feasible. Although there appears to be a surprisingly limited amount of work in the stochastic extensions of the classical Zermelo navigation problem (see [35] for an exception), extensions into the mixed-state domain are desirable for the designing and understanding of more robust controlled quantum dynamics.

Acknowledgments

We thank Eva-Maria Graefe, Benjamin Russell, and Dmitry Savin for discussion and comments, and Martin Meier for support in producing figure 2.

Appendix. Derivation of the universal quantum speed limit

The purpose of this appendix is to offer a proof of the following claim in the paper: *The squared speed of the evolution of a quantum state generated by a Hamiltonian \hat{H} is bounded above by twice the Hilbert–Schmidt norm $\text{tr}(\hat{H}^2)$ of the Hamiltonian, and the bound is attained if and only if \hat{H} is horizontal.*

We begin by establishing some properties of the space of horizontal Hamiltonians. Recall that a Hamiltonian \hat{H} is horizontal with respect to some state $|\psi\rangle$ if and only if $\text{tr}(\hat{H}\hat{H}_{|\psi\rangle}) = 0$ for all Hamiltonians $\hat{H}_{|\psi\rangle}$ that leave $|\psi\rangle$ invariant. For simplicity, we fix the state $|\psi\rangle$ to be the one whose homogeneous coordinates are given by $(1, 0, \dots, 0)^T \in \mathbb{C}^{n+1}$. It should be stressed, however, that the conclusions of the discussion below remain valid for any choice of $|\psi\rangle$, owing to the homogeneous nature of the state space $\mathbb{C}\mathbb{P}^n$. Let us write $V_{|\psi\rangle}$ for the subgroup of $\text{SU}(n+1)$ that leaves $|\psi\rangle$ invariant and $\mathfrak{v}_{|\psi\rangle}$ for the set of the corresponding generators (i.e. its Lie algebra). Then $V_{|\psi\rangle}$ consists of all unit determinant matrices of the form

$$\begin{pmatrix} e^{i\theta} & \mathbf{0}^T \\ \mathbf{0} & \hat{U}_n \end{pmatrix}, \quad \hat{U}_n \in U(n), \quad \mathbf{0}^T = (0, 0, \dots, 0),$$

and $\mathfrak{v}_{|\psi\rangle}$ consists of all trace-free matrices of the form

$$\begin{pmatrix} \lambda & \mathbf{0}^T \\ \mathbf{0} & \hat{B} \end{pmatrix}, \quad \lambda \text{ imaginary and } \hat{B} \in \mathfrak{u}(n).$$

The orthogonal complement $\mathfrak{v}_{|\psi\rangle}^\perp$ of this space with respect to the Hilbert–Schmidt norm coincides, up to a factor of i , with the space of horizontal Hamiltonians. Specifically, the set $\mathfrak{v}_{|\psi\rangle}^\perp$ can be found by requiring that

$$\text{tr} \left(\begin{pmatrix} \lambda & \mathbf{0}^T \\ \mathbf{0} & \hat{B} \end{pmatrix} \begin{pmatrix} \mu & -\bar{\mathbf{z}}^T \\ \mathbf{z} & \hat{C} \end{pmatrix} \right) = \lambda\mu + \text{tr}(\hat{B}\hat{C}) = 0 \quad (9)$$

for all choices of λ and \hat{B} with $\lambda + \text{tr}(\hat{B}) = 0$. Here we have parameterized the full Lie algebra $\mathfrak{su}(n+1)$ using an imaginary μ , a vector $\mathbf{z} \in \mathbb{C}^n$ and $\hat{C} \in \mathfrak{u}(n)$ with $\mu + \text{tr}(\hat{C}) = 0$. It follows that $\mathfrak{v}_{|\psi\rangle}^\perp$ consists of all matrices of the form

$$\begin{pmatrix} 0 & -\bar{\mathbf{z}}^T \\ \mathbf{z} & \mathcal{O} \end{pmatrix},$$

where $\mathbf{z} \in \mathbb{C}^n$ and \mathcal{O} is a null matrix. As a consistency check, we note that this space has $2n$ real dimensions, which of course is the same as the dimensionality of the state space $\mathbb{C}\mathbb{P}^n$. It follows that if \hat{H} is a horizontal Hamiltonian, then

$$2 \text{tr}(\hat{H}^2) = -2 \text{tr} \left(\begin{pmatrix} 0 & -\bar{\mathbf{z}}^T \\ \mathbf{z} & \mathcal{O} \end{pmatrix}^2 \right) = 4 |\mathbf{z}|^2. \quad (10)$$

On the other hand, if $|\psi\rangle$ evolves according to the Schrödinger equation, then the squared speed (in the Fubini–Study metric) of its evolution is given, on account of the Anandan–Aharonov relation, by four times the variance of \hat{H} :

$$4\Delta H^2 = 4 \left(\langle \psi | \hat{H}^2 | \psi \rangle - \langle \psi | \hat{H} | \psi \rangle^2 \right) = 4 |\mathbf{z}|^2. \quad (11)$$

This establishes part of our claim: Namely, by comparing (10) and (11) we see that the square of the evolution speed equals twice the Hilbert–Schmidt norm $\text{tr}(\hat{H}^2)$ if \hat{H} is horizontal. To complete the proof, note that any Hamiltonian \hat{H} can be decomposed into vertical and horizontal parts, i.e. it can be written in the form

$$\hat{H} = i \begin{pmatrix} \lambda & \mathbf{0}^T \\ \mathbf{0} & \hat{B} \end{pmatrix} + i \begin{pmatrix} 0 & -\bar{\mathbf{z}}^T \\ \mathbf{z} & \mathcal{O} \end{pmatrix}. \quad (12)$$

Then

$$2 \text{tr}(\hat{H}^2) = 4 |\mathbf{z}|^2 - 2 \text{tr} \left(\begin{pmatrix} \lambda & \mathbf{0}^T \\ \mathbf{0} & \hat{B} \end{pmatrix}^2 \right), \quad (13)$$

but the trace on the right hand side is strictly negative if the vertical part of \hat{H} is nonzero. Therefore, the squared speed of evolution on state space, given by $4|\mathbf{z}|^2$, is strictly smaller than $2 \text{tr}(\hat{H}^2)$ if the Hamiltonian contains a vertical component. This completes the proof of the claim.

References

- [1] Margolus N and Levitin L B 1998 The maximum speed of dynamical evolution *Physica D* **120** 188
- [2] Lloyd S 1999 Ultimate physical limits to computation *Nature* **406** 1047
- [3] Khaneja N, Glaser S J and Brockett R 2002 Sub-Riemannian geometry and time optimal control of three spin systems: quantum gates and coherence transfer *Phys. Rev. A* **65** 032301
- [4] Palao J P and Kosloff R 2002 Quantum computing by an optimal control algorithm for unitary transformations *Phys. Rev. Lett.* **89** 188301
- [5] Brody D C 2003 Elementary derivation for passage time *J. Phys. A: Math. Gen.* **36** 5587
- [6] Giovannetti V, Lloyd S and Maccone L 2003 Quantum limits to dynamical evolution *Phys. Rev. A* **67** 052109
- [7] Schulte-Herbrüggen T, Spörl A, Khaneja N and Glaser S J 2005 Optimal control-based efficient synthesis of building blocks of quantum algorithms: a perspective from network complexity towards time complexity *Phys. Rev. A* **72** 042331
- [8] Carlini A, Hosoya A, Koike T and Okudaira Y 2006 Time-optimal quantum evolution *Phys. Rev. Lett.* **96** 060503
- [9] Boscaiu U and Mason P 2006 Time minimal trajectories for a spin 1/2 particle in a magnetic field *J. Math. Phys.* **47** 062101
- [10] Brody D C and Hook D W 2006 On optimum Hamiltonians for state transformations *J. Phys. A: Math. Gen.* **39** L167
- [11] Carlini A, Hosoya A, Koike T and Okudaira Y 2007 Time-optimal unitary operations *Phys. Rev. A* **75** 042308
- [12] Rezakhani A T, Kuo W J, Hamma A, Lidar D A and Zanardi P 2009 Quantum adiabatic brachistochrone *Phys. Rev. Lett.* **103** 080502
- [13] Caneva T, Murphy M, Calarco T, Fazio R, Montangero S, Giovannetti V and Santoro G E 2009 Optimal control at the quantum speed limit *Phys. Rev. Lett.* **103** 240501
- [14] Rojo A G and Bloch A M 2010 The rolling sphere, the quantum spin and a simple view of the Landau–Zener problem *Am. J. Phys.* **78** 1014
- [15] Hegerfeldt G C 2013 Driving at the quantum speed limit: optimal control of a two-level system *Phys. Rev. Lett.* **111** 260501
- [16] Garon A, Glaser S J and Sugny D 2013 Time-optimal control of SU(2) quantum operations *Phys. Rev. A* **88** 043422
- [17] Russell B and Stepney S 2014 Applications of Finsler geometry to speed limits to quantum information processing *Int. J. Found. Comput. Sci.* **25** 489
- [18] Wang X, Allegra A, Jacobs K, Lloyd S, Lupo C and Mohseni M 2014 Quantum brachistochrone curves as geodesics: obtaining accurate control protocols for time-optimal quantum gates (arXiv:1408.2465)
- [19] Zermelo E 1931 Über das Navigationsproblem bei ruhender oder veränderlicher Windverteilung *Z. Angew. Math. Mech.* **11** 114
- [20] Carathéodory C 1935 *Variationsrechnung und Partielle Differentialgleichungen erster Ordnung* (Berlin: B G Teubner)
- [21] Russell B and Stepney S 2014 Zermelo navigation and a speed limit to quantum information processing *Phys. Rev. A* **90** 012303
- [22] Russell B and Stepney S 2015 Zermelo navigation in the quantum brachistochrone *J. Phys. A: Math. Theor.* **48** 115303
- [23] Brody D C and Meier D M 2015 Solution to the quantum Zermelo navigation problem *Phys. Rev. Lett.* **114** 100502
- [24] Brody D C and Meier D M 2015 Elementary solution to the time-independent quantum navigation problem *J. Phys. A: Math. Theor.* **48** 055302
- [25] Sunakawa S 1991 *Quantum Mechanics* (Tokyo: Iwanami)
- [26] Bao D, Robles C and Shen Z 2004 Zermelo navigation on Riemannian manifolds *J. Differ. Geom.* **66** 391
- [27] Robles C 2007 Geodesics in Randers spaces of constant curvature *Trans. Am. Math. Soc.* **359** 1633
- [28] Anandan J and Aharonov Y 1990 Geometry of quantum evolution *Phys. Rev. Lett.* **65** 1697
- [29] Kibble T W B 1979 Geometrisation of quantum mechanics *Commun. Math. Phys.* **65** 189
- [30] Bouchiat C and Gibbons G W 1988 Non-integrable quantum phase in the evolution of a spin-1 system: a physical consequence of the non-trivial topology of the quantum state-space. *J. Phys. France* **49** 187
- [31] Brody D C and Hughston L P 2001 Geometric quantum mechanics *J. Geom. Phys.* **38** 19
- [32] Uzdin R, Günther U, Rahav S and Moiseyev N 2012 Time-dependent Hamiltonians with 100% evolution speed efficiency *J. Phys. A: Math. Theor.* **45** 415304
- [33] Kobayashi S and Nomizu K 1969 *Foundations of Differential Geometry* vol 2 (New York: Wiley)
- [34] Gibbons G W and Warnick C M 2011 The geometry of sound rays in a wind *Contemp. Phys.* **52** 197
- [35] Esfahani P M, Chatterjee D and Lygeros J 2011 On a problem of stochastic reach-avoid set characterisation *IEEE Conf. on Decision and Control and European Control Conference* p 7069