

**The Glasgow Voice Memory Test: assessing the ability to memorize and recognize unfamiliar voices**

**Aglieri Virginia<sup>1</sup>, Watson Rebecca<sup>2,3</sup>, Pernet Cyril<sup>4</sup>, Latinus Marianne<sup>1,5</sup>, Garrido Lúcia<sup>6</sup>, Belin Pascal<sup>1,2</sup>**

<sup>1</sup> Institut des Neurosciences de la Timone, UMR 7289, CNRS and Université Aix-Marseille, Marseille, France

<sup>2</sup> Centre for Cognitive Neuroimaging, Institute of Neuroscience and Psychology, University of Glasgow, Glasgow, UK

<sup>3</sup> Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands

<sup>4</sup> Centre for Clinical Brain Sciences, The University of Edinburgh, UK

<sup>5</sup> Unité Mixte de recherche UMR Inserm U930, Université François-Rabelais de Tours, Tours, France

<sup>6</sup> Department of Psychology, Brunel University, London, UK

**Abstract**

**Methods.** 1120 subjects as well as a developmental phonagnosic subject (KH) along with age-matched controls performed the Glasgow Voice Memory Test, which assesses the ability to encode and immediately recognize, through an old/new judgment, both unfamiliar voices (delivered as vowels, making language requirements minimal) and bell sounds. The inclusion of non-vocal stimuli allows the detection of significant dissociations between the two categories (vocal vs non-vocal stimuli).

**Results.** The distributions of accuracy and sensitivity scores ( $d'$ ) reflected a wide range of individual differences in voice recognition performance in the population. As expected, KH showed a dissociation between the recognition of voices and bell sounds, her performance being significantly poorer than matched controls for voices but not for bells.

**Conclusion.** By providing normative data of a large sample and by testing a developmental phonagnosic subject, we demonstrated that the Glasgow Voice Memory Test, available online and accessible from all over the world, can be a valid screening tool (~ 5 min) for a preliminary detection of potential cases of phonagnosia and of “super recognizers” for voices.

### **Introduction**

The ability to recognize familiar faces and match two identical facial configurations between them varies from subject to subject, showing a broad spectrum of individual differences in the normal population. At the lowest extreme of this distribution, there are subjects characterized by an impaired performance in recognizing faces, which have been extensively documented in the literature (Avidan et al., 2014; Avidan, Hasson, Malach, & Behrmann, 2005; Avidan & Behrmann, 2009; Behrmann, Avidan, Gao, & Black, 2007). This deficit, referred to as prosopagnosia, or “face-blindness”, can be present at birth (“developmental phonagnosia”) or acquired after lesions occurring in the ventro-temporal cortex (Barton, 2008). At the opposite extreme, there are individuals with extremely good performance in recognizing faces (“super recognizers”) (Russell, Duchaine, & Nakayama, 2009). To test subjects’ performances, a number of standardized tests are nowadays available such as the Cambridge Face Memory Test (CFMT), which targets the ability to recognize the same face from different points of view and under noisy configurations (e.g. Gaussian noise added to the pictures); therefore, this test recruits a stage of processing which does not require any judgment on the familiarity of the stimuli (Duchaine & Nakayama, 2006). Since its validation in a sample of normal and prosopagnosic subjects, the CFMT has allowed the comparison between different research findings in the domain of face recognition and it has been used to assess individual differences in face recognition (Germine, Duchaine, & Nakayama, 2011; Hedley, Brewer, & Young, 2011).

To date it still remains unclear if the same broad spectrum of performances can be observed in the normal population for the vocal domain. There are evidences that environmental factors contribute to the improvement of the abilities to recognize voices; for instance, an extensive musical training seems to be related to significant higher accuracy in discriminating different voice timbres (Chartrand & Belin, 2006; Chartrand, Peretz, & Belin, 2008). Furthermore, cases of developmental phonagnosia have been recently described, pointing out that in the general population there could be a specific deficit for the recognition of vocal stimuli which does not result from any neurological lesion (Garrido et al., 2009; Herald, Xu, Biederman, Amir, & Shilowich, 2014; Roswadowitz et al., 2014). Developmental phonagnosia can be viewed as the equivalent of developmental prosopagnosia in the vocal domain and its investigation is fundamental to better understand models of person-recognition, particularly in the light of recent findings of multisensory integration of facial and vocal cues in person-recognition processes (von Kriegstein et al., 2008; von Kriegstein, Kleinschmidt, & Giraud, 2006). Similarly to prosopagnosia, acquired phonagnosia can be observed either for familiar voices (D. R. Van Lancker, Kreiman, & Cummings, 1989; D. R. Van Lancker & Canter, 1982) or non-familiar voices (Jones et al, in revision ) in patients with specific lesions of the right parietal vs. right inferior frontal cortices.

Despite these known deficits, there is no agreement on which tests to use to reliably detect and document voice deficits. Indeed, no test validation in phonagnosic and normal subjects has been performed so far. The tests used in previous research on vocal processing were usually created for the purpose of the study and, often, dependent on the language of participants. Furthermore, studies investigating acquired phonagnosia in brain-lesioned patients used both discrimination and recognition tasks (Hailstone, Crutch, Vestergaard, Patterson, & Warren, 2010; Neuner & Schweinberger, 2000; D. R. Van Lancker, Cummings,

Kreiman, & Dobkin, 1988). Since recognizing familiar voices and discriminating unfamiliar ones seem to involve different areas in the brain (D. Van Lancker & Kreiman, 1987), it remains impossible to unequivocally associate a lesion site with acquired phonagnosia because the data gathered so far are based on patients tested with these two different types of tests.

Given the need for standardization and reproducibility in the field of voice processing, we here present the Glasgow Voice Memory Test validated in a sample of 1120 subjects gathered online in comparison with the first published case of developmental phonagnosia, KH (Garrido et al., 2009). This brief test (5 minutes) targets perceptual and memory aspects of vocal processing by comparing the performance obtained in encoding both vocal stimuli and bell sounds and immediately judging the stimuli as familiar or unfamiliar. This allows us to evaluate performance level at voice encoding and familiarity recognition, and look at potentially significant dissociations between the vocal and non-vocal domains (Crawford & Garthwaite, 2005). The inclusion of the same task repeated for both voices and bell sounds is in line with the idea behind the development of the Cambridge Car Memory Test (CCMT; Dennett et al., 2012), which requires to learn and recognize cars with the same procedure used in the Cambridge Face Memory Test. Cars, as bells, are stimuli that allow to investigate the ability to discriminate different examples within an object category. According to the data gathered in a large sample of subjects, the CFMT and the CCMT seem to tap into different processes (Dennett et al., 2012).

The GVMT is currently available online (<http://experiments.psy.gla.ac.uk/>) and, hence, easily accessible from all over the world. One of its main strengths is that of presenting vocal stimuli characterized by minimal verbal information (the vowel /a/), which makes it an optimal tool not only for comparing the performance of subjects of different

nationalities, but also to be used (in a not online version) in all kind of neurological patients, including aphasic ones.

By analyzing the data gathered online from a large and heterogeneous sample of subjects, we expected to observe a wide range of individual differences in voice recognition abilities, as has been observed for faces. Furthermore, we hypothesized that the developmental phonagnosic subject KH would show a significant poorer performance compared to matched controls in voice recognition but not in the recognition of bells, demonstrating the validity of the GVMT. Finally, norms are presented in the appendix allowing to compare any new subject to our sample.

## **Methods**

### *Online test*

1120 adults aged 18 upwards performed the test online (743 females;  $M=26.7$  years,  $SD = 11.1$ , range [18-86]). There were in total 59 different nationalities. In order to take part to the experiment, it was required to first register to the website by giving informed consent. Participants were asked to indicate their age, if they had a twin (in this case, to provide his/her email) and to self-assess their hearing abilities (normal, impaired or presence of hearing deficits such as tinnitus). Only participants that stated to have normal hearing abilities were included in the test. The instructions for the experiment were then displayed (“Your task is to listen to a series of eight voices and try to remember them. This will be followed by another series of voices that will test your memory. For each one of those new voices, you will have to indicate if it belongs to the first series you have been trying to remember. This will be repeated for ringing bells”). A sound test was made available in order to try if the speakers of the device used were correctly operating. Upon completion, participants were given their own score as well as an indication of how well they performed compared to the

general population (in percentage). The study was approved by the local ethics committee, and was run according to the Helsinki guidelines.

### *Phonagnosic subject (KH) and controls*

KH is a right handed woman aged 62 at the time of testing, who reported to be unable to recognize voices of famous people and of her friends and family. Her case has been fully described in Garrido et al., (2009). She was tested against a control group composed of 6 women matched for age (M=58 years, range [52-68]) and relative level of education. The participation of KH was on a voluntary basis. The participants of the control group were rewarded at the usual rate paid by University of Glasgow (£6 per hour).

### *Stimuli*

A total of 16 voices (8 male) with a mean duration of 487 ms and the recorded sounds of 16 different bells of mean duration of 1110 ms were used. Voice stimuli (French vowel /a/) were obtained from recordings performed in Montreal. The native language of all speakers was Canadian French. Recordings (16 bit, 44.1kHz) of the speakers were made in the multi-channel recording studio of Secteur ElectroAcoustique in the Faculté de musique, Université de Montreal, using two Bruel & Kjaer 4006 microphones (Bruel & Kjaer; Nærum, Denmark), a Digidesign 888/24 analog/digital converter and the Pro Tools 6.4 recording software (both Avid Technology; Tewksbury, MA, USA). Bell sounds were obtained from a public internet source ([www.findsounds.com](http://www.findsounds.com)).

### *Procedure*

The test was structured into four phases: 1) encoding of voices; 2) recognition of voices; 3) encoding of bells; 4) recognition of bells.

- 1) Encoding of voices

Participants initially heard 8 voices. The first four voices delivered were of females, while the other four of males. Each voice was presented 3 times in a row, with an interstimulus interval (ISI) between the onsets of the sounds of 1500 ms; different triplets were separated by a 3000 ms silent gap. The presentation order during the encoding phase was the same for all subjects.

## 2) Recognition of voices

After the encoding phase ended, participants were asked to start the recognition phase whenever they were ready, while another sound-check was made available. During this phase, participants heard the 8 voices presented during the encoding phase and 8 new ones (4 of females and 4 of males). Voices were presented in a random order. Subjects performed an old/new task on the stimuli: they had to decide whether the voice they heard had been presented in the encoding phase ('old') or if had not been presented ('new'). The decision was self-paced. Between participants' decision and the loading of the next sound there was an interval of 1000 ms.

## 3) Encoding of bells

During this phase, participants were instructed to listen to 8 different sounds of bells. The presentation procedure was the same as for the vocal stimuli.

## 4) Recognition of bells

After the encoding phase for bells ended, participants were asked to start the recognition phase. During this phase, participants heard the 8 bells presented during the encoding phase and 8 new ones. Bells were here presented in a random order. Subjects performed an old/new task on the stimuli: they had to decide whether each voice had been presented in the encoding phase ('old') or not ('new'). The decision was self-paced. Between participant's decision and the loading of the next sound there was an interval of 1000ms.

Thus, instructions delivered and task demands were highly similar for the voice and the bells part of the GVMT.

*Data analysis*

For both tasks, we analyzed data in line with detection theory (Macmillan, 2002; Macmillan & Creelman, 2004), measuring hit rates (HR; a voice previously heard was correctly classified as old), false alarms (FA; a voice heard for the first time was classified as old), misses (an old voice considered new) and correct rejections (CR; a voice never heard was classified as new). We calculated the percent correct (PC), which takes into account both hit rates and correct rejections ( $PC = ((HR + CR) / 2) * 100$ ), and  $d'$  ( $d$  prime), computed instead as the difference between standardized hit rates and false alarms. Hence, percent correct is a measure indicative of both sensitivity (proportion of actual positives correctly identified as such) and specificity (proportion of negatives correctly identified as such), while  $d'$  is used as a measure of participants' sensitivity to correctly identify a previously heard stimulus as old.

All statistical analyses applied to compare KH's performance to matched controls followed the guidelines provide in Crawford & Howell (1998). The modified t-test is adapted for comparing one single case to a small group of control subjects. Furthermore, when testing a patient, it is important to show a significant dissociation between the performances obtained in two different tasks, likely tapping into different cognitive and neural processes. To test if KH was impaired in recognition of voices but not of bells, we ran a revised standardized difference test for dissociations (Crawford & Garthwaite, 2005). When needed, robust skipped correlations (Spearman) were computed to protect against the effect of marginal and bivariate outliers (Pernet, Wilcox, & Rousselet, 2013). In this method, the acceptance or rejection of the null hypothesis is performed on bootstrap 95 % confidence intervals to protect against heteroscedasticity (e.g. if the CIs do not include 0, the null hypothesis of no correlation can be refused).



All the analyses were run in MATLAB (MATHWORKS Inc., Natick, MA) using statistical toolbox.

## Results

The distributions of the scores of the 1120 subjects calculated as percent correct (PC) and  $d'$  are showed in Fig. 1 (boxplots) & Fig. 2 (histograms). The Jarque-Bera test, which tests the null hypothesis that the data set has skewness and kurtosis matching a normal distribution (hence both these measures being equal to zero) (Gel & Gastwirth, 2008), revealed violation of normality for both percent correct and  $d'$  scores, for both voices and bells (all  $p < 0.001$ ). More specifically, the distributions were all negatively skewed, having most of the scores clustered on the right (higher performance levels); this violation of skewness could indicate a ceiling effect. Referring to kurtosis values ( $k$ ), the distribution of percent correct for voices was platykurtic ( $k < 0$ ), having a peak lower and broader than expected for normally distributed values, while for bell recognition was leptokurtic ( $k > 0$ ), having a central peak higher and sharper. The distributions for  $d'$  scores for voices and bells were instead both platykurtic.

Since it is possible that a bad performance in voice recognition is accompanied by a comparable bad performance in recognition for bells, we also looked at the distribution of the differences between the two performances (voice – bells), which allows to focus on significant dissociations. This distribution (PC for voices – PC for bells;  $M = -5.24$ ;  $SD = 12.82$ ;  $CI (95\%) = [-5.99, -4.49]$ ) was normal (Jarque-Bera test,  $p = 0.3$ ). The difference between  $d'$  scores for voices and bells ( $M = -0.34$ ;  $SD = 0.8268$ ;  $CI (95\%) = [-0.39, 0.29]$ ) also followed a normal distribution (Jarque-Bera test;  $p = 0.21$ ) (Fig. 3 & 4).

Since both mean differences were negative, we assessed through a Wilcoxon matched-pair test if bells sounds were significantly better recognized than voices. The results show that this was the case for both percent correct ( $Z = -12.87$ ,  $p < 0.001$ , effect size:  $r =$

0.27) and  $d'$  ( $Z = -12.69$ ,  $p < 0.001$ , effect size:  $r = 0.28$ ). Nevertheless, there was a significant positive correlation between the performance for voices and bells, both for percent correct scores (skipped Spearman correlation;  $\rho = 0.2$ ,  $t = 6.98$ ,  $CI (95\%) = [0.14, 0.26]$ ), and  $d'$  scores (skipped Spearman correlation;  $\rho = 0.21$ ,  $t = 7.33$ ,  $CI (95\%) = [0.16, 0.27]$ ).

To investigate possible gender effects on the general performance, we compared the differences between males' and females' performances in voice and bell recognition with a  $t$ -test assuming unequal variances. The results point out to a null effect of gender of the listener, both for  $d'$  ( $t (747.2865) = -0.122$ ,  $CI (95\%) = [-0.11, 0.1]$ ,  $p > 0.05$ ) and percent correct ( $t (754.2623) = -0.07$ ,  $CI (95\%) = [-1.65, 1.53]$ ,  $p > 0.05$ ) (Fig. 5).

No significant correlation was found between PC scores for voices and age of participants (Skipped Spearman correlation;  $\rho = 4.0132e-04$ ,  $t = 0.0134$ ,  $CI (95\%) = [-0.07, 0.06]$ ) nor between PC scores for bells and age (Skipped Spearman correlation;  $\rho = 8.0259e-04$ ,  $t = 0.027$ ,  $CI (95\%) = [-0.06, 0.06]$ ). The same pattern was also observed for  $d'$  scores for voices (Skipped Spearman correlation;  $\rho = 0.0087$ ,  $t = 0.29$ ,  $CI (95\%) = [-0.05, 0.07]$ ) and for bells ( $\rho = -0.014$ ,  $t = -0.47$ ,  $CI (95\%) = [-0.078, 0.053]$ ).

Appendix 1 provides the detailed distributions of all the measures of interest by percentiles.

	Observed range		All (N=1120)			Females (N=743)			Males (N=377)		
	Min	Max	M	SD	95 % CI	M	SD	95 % CI	M	SD	95 % CI
<b>Age</b>	18	86	26.7	11.10	[26, 27.3]	25.89	10.47	[25.13, 26.64]	28.17	12.11	[28.17, 26.94]
<b>PC voices (%)</b>	37.5	100	78.15	10.95	[77.5, 78.79]	77.89	10.75	[77.12, 78.67]	78.65	11.33	[77.5, 79.79]
<b>D' voices</b>	-0.67	3.07	1.66	0.69	[1.61, 1.7]	1.64	0.68	[1.59, 1.69]	1.69	0.72	[1.61, 1.76]
<b>PC bells (%)</b>	43.75	100	83.39	9.97	[82.81, 83.98]	83.85	9.77	[82.45, 83.86]	83.16	10.33	[82.8, 84.9]
<b>D' bells</b>	-0.35	3.07	1.99	0.64	[1.95, 2.03]	1.98	0.63	[1.93, 2.02]	2.02	0.66	[1.95, 2.09]
<b>PC voices – PC bells</b>	-43.75	43.75	-5.24	12.82	[-5.99, -4.49]	-5.26	12.82	[-6.19, -4.34]	-5.21	12.85	[-6.51, -3.9]

<b>d'</b> <b>voices –</b> <b>d' bells</b>	-2.74	2.56	-0.34	0.83	[-0.39 ,0.29]	-0.34	0.82	[-0.4, -0.28]	-0.33	0.83	[-0.42 ,0.25]
---	-------	------	-------	------	------------------	-------	------	---------------	-------	------	------------------


**Table 1. Summary statistics of the online sample.** Range, means, standard deviations (SD) and 95 % confidence intervals observed for age and scores obtained in voices and bells recognition and their differences (PC= percent correct; d'=d primes).

KH's percent correct scores were significantly lower than those of age-matched controls for voice recognition ( $t(5) = -2.04$ ;  $p = 0.05$ ; effect size = -2.2) but not for bell recognition ( $t(5) = 1.19$ ;  $p > 0.05$ ; effect size = 1.29) (see Table 2), as confirmed by the result of the revised standardized difference test for dissociations ( $t(5) = 2.85$ ,  $p = 0.018$ ). D' for voices was significantly smaller for KH than for controls ( $t(5) = -2.04$ ,  $p < 0.05$ ; effect size = -2.2); d' for bell recognition did not differ between KH and controls ( $|t(5)| < 1.5$ ; effect size = 1.33). The revised standardized difference test for dissociations confirmed a significant dissociation in KH also for d' scores ( $t(5) = 2.71$ ,  $p = 0.02$ ) (Fig. 6).


	Voices		Bells	
	PC	d'	PC	d'
<b>KH</b>	50	0	93.75	2.68
<b>Controls (N=6)</b>	72 ± 10	1.32 ± 0.6	77 ± 13	1.64 ± 0.78
<b><i>t(5)</i></b>	<b>-2.04*</b>	<b>-2.04*</b>	1.19	1.23
<b>Effect size on <i>t(5)</i> [95 % CI]</b>	-2.2 [-3.72, -0.64]	-2.2 [-3.18, 0.63]	1.29 [0.14, 2.37]	1.33 [0.17, 2.44]

**Table 2. Mean and SDs of percent correct and d' scores and results of the modified t-test (*t(5)*) comparing KH's performance and matched controls.** The negative values indicate that KH's performance was significantly poorer than for controls. The third column reports the results of the revised t-test for differences between a single case and controls. Values presented in bold are significant (one-tailed,  $p < 0.05$ ).


The effect size is reported together with its relative confidence interval.




**Figure 1. Boxplots representing performance distribution.** Percent correct scores (left) and  $d'$  scores (right) for recognition of voices and bells. Red crosses represent scores corresponding to 2 SDs below or above the average.  $N=1120$ .




**Figure 2. Histograms representing the distribution of performances.** Percent correct scores (top) and  $d'$  scores (bottom) for recognition of voices (left) and bells (right). The red asterisk indicates the performance obtained by the phonagnosic subject KH overlaid on the results of the 1120 subjects of the online test.




**Figure 3. Boxplots for the distribution of the differences between performances.** Differences between the two performances (voice recognition – bell recognition) for PC (left) and  $d'$  (right) scores in 1120 subjects. Red crosses represent scores 2 SDs below or above the average.



**Figure 4. Distribution of the differences between the two performances (voice recognition – bell recognition) for both PC (left) and  $d'$  (right) scores.** Red asterisks indicate KH's performance overlaid on the results of the 1120 subjects of the online test.



**Figure 5. Bar graphs representing mean PC and d' scores for recognition of voices and bells, separated by gender of the listener. Error bars represent 95% confidence intervals.**



**Figure 6. Bar graphs representing PC (left) and d' scores (right) of KH and matched controls. A dissociation was observed between the recognition of voices and bell sounds. KH's performance was significantly poorer than that of controls for recognition of voices but not of bells. Error bars represent 95% confidence intervals (\*p<0.05).**

When investigating individual differences, it is also advisable to compute measures of inter-subject reliability; this type of measure can in fact inform us on whether the participants classified voices and bells in a consistent way among them. Hence, we analyzed the dichotomous variable of choice (old or new voice/bell) of each participant for each voice (or

bell) using a two-way random effects intra-class correlation (ICC) model. Since we had access to this dichotomous variable of only one part of our subjects ( $N = 598$ ), these analysis did not include the entire sample. The partial results point out to a fair agreement among 598 raters in the classification of voices (ICC coefficient = 0.38, CI (95%) = [0.25, 0.6],  $F(15) = 373.89$ ) and a moderate agreement in the classification of bells (ICC coefficient = 0.52, CI (95%) = [0.37, 0.72],  $F(15) = 645$ ).

Furthermore, testing internal consistency is fundamental for assessing that the different items of a test target the same construct (e.g. different voices all testing the ability to recognize voices). For this purpose, we also checked the internal consistency of the GVMT by looking again at the dichotomous variable of choice (old or new) for both categories of stimuli. The results point out to an optimal internal consistency of our test, for both voices (Cronbach's alpha = 0.9973) and bells (Cronbach's alpha = 0.9984). These coefficients have also been computed on a smaller sample of subjects ( $N = 598$ ).

## Discussion

We here summarize the major results gathered in a sample of subjects that performed the GVMT online as well as in a developmental phonagnosic subject (KH) and matched controls.

*GVMT: a tool for investigating individual differences in voice processing abilities*

The normative data obtained in a sample of 1120 subjects of different ages and cultures highlights a wide range of individual differences in the ability to encode and immediately recognize unfamiliar voices. Interestingly, the distributions of the differences for both PC and  $d'$  showed that there were cases in which an extremely poor performance in voice recognition was accompanied by an extremely good performance in recognition of bells, meaning that this pattern cannot be ascribed to a general deficit in auditory processes or to difficulties posed by the task. It has been previously demonstrated that the contrast

between vocal and environmental stimuli lead to the activation of specific areas in the temporal lobe and superior temporal sulcus, named the Temporal Voice Areas (TVAs; (Belin, Zatorre, Lafaille, Ahad, & Pike, 2000). Furthermore, the functional activity in the TVAs during passive listening of sounds compared to baseline (vocal + non vocal sounds >baseline) was found to predict the performance for voice recognition obtained in the GVMT (Watson, Latinus, Bestelmeyer, Crabbe, & Belin, 2012). Hence, future studies should look at the functional activity in these areas while the GMVT is performed in order to associate individual differences in behavior to different patterns of neural activity.

According to the results of the inter-subject reliability analysis, it seems that there is slightly more variability in the way subjects classified the 16 vocal stimuli presented than the 16 environmental ones; this tendency points out to the fact that the subjective saliency attributed to vocal stimuli could vary more among subjects than the one attributed to other stimuli.

*GVMT: a reliable and valid screening test for the detection of phonagnosia*

In terms of reliability (the degree to which an assessment tool produces stable and consistent results), our results show that the GVMT has optimal internal consistency reliability, meaning that the different items chosen (e.g. the 16 different voices and the 16 environmental stimuli) consistently test the same construct.

The GVMT seems also to be a valid test for the assessment of voice recognition abilities because KH, the first documented case of developmental phonagnosia (Garrido et al., 2009), presented a dissociation between recognition of voices and bells. She performed significantly worse than matched controls in voice recognition but better in the recognition of bells (even if this difference did not reach significance). Although there are no formal criteria available to declare a subject as phonagnosic, the extensive assessment performed on KH in 2009 seemed to point out to the presence of a deficit in recognizing and discriminating voices



in presence of intact auditory abilities and general sound processing. Garrido et al., (2009) observed in fact that KH was impaired in both recognition of voices of celebrities and discrimination of different vocal stimuli but that she was as good as matched controls in recognizing environmental sounds and in processing musical stimuli. Here, even a simple task such as an old/new judgment on voices and bells heard for the first time lead to similar results.

Since the GVMT seems to specifically detect a deficit in vocal processing, we propose that it could be used as an initial screening tool in the assessment of this deficit in both the general population (to investigate developmental phonagnosia) and neurological patients (to investigate acquired phonagnosia). It is advisable, in any case, that a more extensive assessment tapping into higher stages of processing such as identity recognition as the one used by Garrido et al., (2009) and more recently by Roswadowitz et al., (2014) is also carried out to detect a specific impairment in the recognition of voices. To date, we cannot in fact confirm that the GVMT is sensitive to different types of phonagnosia. There seems in fact to exist an apperceptive form of phonagnosia, resulting in an impaired performance in perceptual matching tasks, and an associative phonagnosia, which refers to the inability to associate semantic information to a voice (Roswadowitz et al., 2014). According to the results in Roswadowitz et al., (2014), a subject with apperceptive phonagnosia could be detected through a discrimination task which requires to perform a judgment of similarity between two voices; at the contrary, a subject with associative phonagnosia could present a spared performance in a discrimination task but would be significantly impaired in a test that requires to provide semantic information associated to the voice of a famous or personally known person. By looking at the performance of KH in the GVMT, it is not clear to which type of phonagnosia KH belongs; the test here presented, in fact, does not specifically assess

voice discrimination or recognition. Rather, it tests the ability to activate a sense of familiarity toward a stimulus briefly presented for the first time.

### *Limitations*

One of the criticisms that might be raised to the GVMT is that it taps more into short-term memory abilities than specific abilities to process vocal sounds. Nevertheless, if this was the case, significant dissociations between recognition of voices and bells such as in KH and in other subjects that performed the test online should not be observed.

According to our results, environmental sounds such as bells seem to be easier to recognize than voices. This finding should be carefully considered since it cannot be excluded that there was an order effect; the test for bells was in fact always presented after the test for voices, when subject already familiarized with the procedure. Furthermore, the bell stimuli here used lasted longer than vocal ones, and it has been shown that voice recognition improves with increasing duration of vocal samples (D. Van Lancker, Kreiman, & Emmorey, 1985).

Another limitation of our study (and, in general, of online testing) is that we discarded the analysis of reaction times because they could be affected by different speeds of internet connections and operating systems and by the fact that subjects are not controlled by the experimenter; hence, we do not have any information on possible differences in processing time of the two types of stimuli, which would instead be useful to compute measures of speed/accuracy trade off, as previously done in prosopagnosic subjects (Busigny, Joubert, Felician, Ceccaldi, & Rossion, 2010). Furthermore, we could not control for the time occurred between the encoding and recognition phases; even if it is more likely that, being the test particularly short, participants completed it without taking long breaks, it cannot be excluded that this interval considerably varies among subjects. Nevertheless, these limitations

related to the timing of experiment could be overcome by comparing performance on the GVMT online and in the laboratory.

Despite these limitations, a web-based experiment such as the one here presented can have a great potential in identifying cases of phonagnosia in the general population as it allows for the gathering of large samples of data, overcoming issues related to small sample sizes.

### Acknowledgements

This work was supported by grants BB/E003958/1 from BBSRC, large grant RES-060-25-0010 by ESRC/MRC, and grant AJE201214 by the Fondation pour la Recherche Medicale.

VA is supported by a PhD fellowship from the A\*MIDEX foundation. We are grateful to Guylaine Bélizaire, Maude Urfer, Cyril Pernet and Marc Becirspahic for their contribution to developing earlier versions of this test. We do not declare any conflict of interest.

### References

- Avidan, G., & Behrmann, M. (2009). Functional MRI Reveals Compromised Neural Integrity of the Face Processing Network in Congenital Prosopagnosia. *Current Biology*, *19*(13), 1146–1150.
- Avidan, G., Hasson, U., Malach, R., & Behrmann, M. (2005). Detailed Exploration of Face-related Processing in Congenital Prosopagnosia: 2. Functional Neuroimaging Findings. *Journal of Cognitive Neuroscience*, *17*(7), 1150–1167.
- Avidan, G., Tanzer, M., Hadj-Bouziane, F., Liu, N., Ungerleider, L. G., & Behrmann, M. (2014). Selective dissociation between core and extended regions of the face processing network in congenital prosopagnosia. *Cerebral Cortex*, *24*(6), 1565–1578.
- Barton, J. J. (2008). Structure and function in acquired prosopagnosia: lessons from a series of 10 patients with brain damage. *Journal of Neuropsychology*, *2*(1), 197–225.

- Behrmann, M., Avidan, G., Gao, F., & Black, S. (2007). Structural Imaging Reveals Anatomical Alterations in Inferotemporal Cortex in Congenital Prosopagnosia. *Cerebral Cortex*, *17*(10), 2354–2363.
- Belin, P., Zatorre, R. J., Lafaille, P., Ahad, P., & Pike, B. (2000). Voice-selective areas in human auditory cortex. *Nature*, *403*(6767), 309–312.
- Busigny, T., Joubert, S., Felician, O., Ceccaldi, M., & Rossion, B. (2010). Holistic perception of the individual face is specific and necessary: Evidence from an extensive case study of acquired prosopagnosia. *Neuropsychologia*, *48*(14), 4057–4092.
- Chartrand, J.-P., & Belin, P. (2006). Superior voice timbre processing in musicians. *Neuroscience Letters*, *405*(3), 164–167.
- Chartrand, J.-P., Peretz, I., & Belin, P. (2008). Auditory recognition expertise and domain specificity. *Brain Research*, *1220*, 191–198.
- Crawford, J. R., & Garthwaite, P. H. (2005). Testing for suspected impairments and dissociations in single-case studies in neuropsychology: evaluation of alternatives using monte carlo simulations and revised tests for dissociations. *Neuropsychology*, *19*(3), 318.
- Crawford, J. R., & Howell, D. C. (1998). Comparing an individual's test score against norms derived from small samples. *The Clinical Neuropsychologist*, *12*(4), 482–486.
- Dennett, H. W., McKone, E., Tavashmi, R., Hall, A., Pidcock, M., Edwards, M., & Duchaine, B. (2012). The Cambridge Car Memory Test: A task matched in format to the Cambridge Face Memory Test, with norms, reliability, sex differences, dissociations from face memory, and expertise effects. *Behavior Research Methods*, *44*(2), 587–605.

- Duchaine, B., & Nakayama, K. (2006). The Cambridge Face Memory Test: Results for neurologically intact individuals and an investigation of its validity using inverted face stimuli and prosopagnosic participants. *Neuropsychologia*, *44*(4), 576–585.
- Garrido, L., Eisner, F., McGettigan, C., Stewart, L., Sauter, D., Hanley, J. R., ... Duchaine, B. (2009). Developmental phonagnosia: a selective deficit of vocal identity recognition. *Neuropsychologia*, *47*(1), 123–131.
- Gel, Y. R., & Gastwirth, J. L. (2008). A robust modification of the Jarque–Bera test of normality. *Economics Letters*, *99*(1), 30–32.
- Germine, L. T., Duchaine, B., & Nakayama, K. (2011). Where cognitive development and aging meet: Face learning ability peaks after age 30. *Cognition*, *118*(2), 201–210.
- Hailstone, J. C., Crutch, S. J., Vestergaard, M. D., Patterson, R. D., & Warren, J. D. (2010). Progressive associative phonagnosia: a neuropsychological analysis. *Neuropsychologia*, *48*(4), 1104–1114.
- Hedley, D., Brewer, N., & Young, R. (2011). Face recognition performance of individuals with Asperger syndrome on the Cambridge Face Memory Test. *Autism Research*, *4*(6), 449–455.
- Herald, S. B., Xu, X., Biederman, I., Amir, O., & Shilowich, B. E. (2014). Phonagnosia: A voice homologue to prosopagnosia. *Visual Cognition*, (ahead-of-print), 1–3.
- Jones, A.B., Farrall, A., Belin, P. & Pernet, C.R (2015). Hemispheric association and dissociation of voice and speech information processing in stroke. *Cortex*, in revision.
- Lancker, D. R. V., Kreiman, J., & Cummings, J. (1989). Voice perception deficits: Neuroanatomical correlates of phonagnosia. *Journal of Clinical and Experimental Neuropsychology*, *11*(5), 665–674.
- Macmillan, N. A. (2002). Signal detection theory. *Stevens' Handbook of Experimental Psychology*.

- Macmillan, N. A., & Creelman, C. D. (2004). *Detection theory: A user's guide*. Psychology press.
- Neuner, F., & Schweinberger, S. R. (2000). Neuropsychological impairments in the recognition of faces, voices, and personal names. *Brain and Cognition*, *44*(3), 342–366.
- Pernet, C. R., Wilcox, R. R., & Rousselet, G. A. (2013). Robust correlation analyses: false positive and power validation using a new open source Matlab toolbox. *Quantitative Psychology and Measurement*, *3*, 606.
- Roswadowitz, C., Mathias, S. R., Hintz, F., Kreitewolf, J., Schelinski, S., & von Kriegstein, K. (2014). Two Cases of Selective Developmental Voice-Recognition Impairments. *Current Biology*.
- Russell, R., Duchaine, B., & Nakayama, K. (2009). Super-recognizers: People with extraordinary face recognition ability. *Psychonomic Bulletin & Review*, *16*(2), 252–257.
- Van Lancker, D., & Kreiman, J. (1987). Voice discrimination and recognition are separate abilities. *Neuropsychologia*, *25*(5), 829–834.
- Van Lancker, D., Kreiman, J., & Emmorey, K. (1985). Familiar voice recognition: Patterns and parameters. Part I: Recognition of backward voices. *Journal of Phonetics*, *13*(1), 19–38.
- Van Lancker, D. R., & Canter, G. J. (1982). Impairment of voice and face recognition in patients with hemispheric damage. *Brain and Cognition*, *1*(2), 185–195.
- Van Lancker, D. R., Cummings, J. L., Kreiman, J., & Dobkin, B. H. (1988). Phonagnosia: A Dissociation Between Familiar and Unfamiliar Voices. *Cortex*, *24*(2), 195–209.

- von Kriegstein, K., Kleinschmidt, A., & Giraud, A.-L. (2006). Voice recognition and cross-modal responses to familiar speakers' voices in prosopagnosia. *Cerebral Cortex*, *16*(9), 1314–1322.
- von Kriegstein, K. von, Dogan, Ö., Grüter, M., Giraud, A.-L., Kell, C. A., Grüter, T., ... Kiebel, S. J. (2008). Simulation of talking faces in the human brain improves auditory speech recognition. *Proceedings of the National Academy of Sciences*, *105*(18), 6747–6752.
- Watson, R., Latinus, M., Bestelmeyer, P. E., Crabbe, F., & Belin, P. (2012). Sound-induced activity in voice-sensitive cortex predicts voice memory ability. *Frontiers in Psychology*, *3*.

### Appendix I

Percent Correct	Voices	Bells
0	0	0
6,25	0	0
12,5	0	0
18,75	0	0
25	0	0
31,25	0	0
37,5	0,089	0
43,75	0,18	0,27
50	1,96	0,45
56,25	5,535	1,52
62,5	12,95	4,82
68,75	26,34	12,05
75	48,21	28,84
81,25	69,01	49,73
87,5	87,77	74,55
93,75	97,59	93,48
100	100	100

**Table 1. Quantiles for PC for voice and bell recognition.** The first column reports possible scores divided in 17 intervals, while the other two the percentage of subjects that obtained the corresponding equal or lower score in voice and bell recognition ( $N$  subjects with = or < score / 1120).

d' Score	Voices	Bells
0	1.96	0.45
0.38	5.09	1.43
0.57	5.53	1.52
0.77	12.05	4.2
0.96	12.95	4.82
1.15	25.45	11.78
1.34	26.34	12.05
1.73	48.21	28.84
2.11	69.02	49.73
2.30	87.77	74.55
2.49	87.77	74.55



<b>2.68</b>	97.59	93.48
<b>2.88</b>	97.59	93.48
<b>3.07</b>	100	100

**Table 2. Quantiles for  $d'$  for voice and bell recognition.** The first column reports possible scores divided in 14 intervals, while the other two the percentage of subjects that obtained the corresponding equal or lower score in voice and bell recognition.

<b>PC voices – PC bells</b>	<b>Quantile</b>
<b>-31.25</b>	2.95
<b>-25</b>	9.73
<b>-18.75</b>	20.71
<b>-12.5</b>	37.77
<b>-6.25</b>	56.87
<b>0</b>	74.37
<b>6.25</b>	87.86
<b>12.5</b>	95.18
<b>18.75</b>	98.30
<b>25</b>	99.2
<b>31.25</b>	99.82
<b>37.5</b>	99.91
<b>43.75</b>	100

**Table 3. Quantiles for PC differences between voices and bells.** The first column reports possible scores divided in 13 intervals, while the other the percentage of subjects that obtained the corresponding equal or lower score.

<b>D' voices – d' bells</b>	<b>Quantile</b>
<b>-2.08</b>	1.07
<b>-1.75</b>	2.95
<b>-1.42</b>	9.37
<b>-1.09</b>	19.46
<b>-0.76</b>	34.64
<b>-0.43</b>	45.62
<b>-0.1</b>	58.03
<b>0.23</b>	74.46
<b>0.56</b>	87.86
<b>0.89</b>	94.2
<b>1.22</b>	96.96
<b>1.55</b>	98.48
<b>1.88</b>	99.37
<b>2.21</b>	99.91
<b>2.87</b>	100

**Table 4. Quantiles for d' differences between voices and bells.** The first column reports possible scores divided in 15 intervals, while the other the percentage of subjects that obtained the corresponding equal or lower score.