

*Article*

Running Head: MUSIC AND MUSIC-VIDEO

Bird, J. M., Hall, J., Arnold, R., Karageorghis, C. I., & Hussein, A. (2016). Effects of music and music-video on core affect during exercise at the lactate threshold. *Psychology of Music*. Advance online publication. doi:10.1177/0305735616637909

**Effects of music and music-video on core affect during exercise at the lactate threshold**

Resubmitted: December 23, 2015

**Abstract**

The objective of the present study was to examine the effects of music and music-video on core affect during and immediately after stationary cycling at the lactate threshold. A randomized, fully counterbalanced, crossover design with three conditions (music, music-video, and a no-music-video control) was adopted. Twenty-four participants exercised at lactate threshold while exposed to music, music-video, and control conditions. Affective valence and perceived activation were assessed every 2 min during a 20-min exercise bout and every 5 min post-exercise over a 20-min period. Results indicated that there was a significant condition x time interaction for affective valence. The music-video condition elicited the highest levels of affective valence followed by the music condition and control. There was a main effect of condition for affective valence, wherein the experimental conditions facilitated significantly higher affective valence than control. Significant main effects of time emerged for both affective valence and perceived activation. Regardless of condition, affective valence decreased during the exercise bout and increased immediately post-exercise. Conversely, perceived activation increased during exercise and decreased immediately post-exercise. The present findings indicate that music and music-video can enhance core affect during exercise at the lactate threshold and the implications for exercise adherence are expounded.

**Keywords**

Audio-visual aids, circumplex model, musicology, physical activity, public health

Regular physical activity has been linked to numerous physical and mental health benefits (Chief Medical Officers of England, Scotland, Wales, & Northern Ireland, 2011). Despite such benefits, the percentage of people engaging in regular physical activity has remained low over the last two decades (Ekkekakis & Dafermos, 2012), to the extent that its promotion has become a “key public health issue” in the UK (Biddle et al., 2010, p. 5). Using hedonic theory (Kahneman, 1999) as a guiding framework, researchers have emphasized the role of affective responses to exercise as determinants of exercise adherence (Tempest & Parfitt, 2013; Williams, 2008), suggesting that human behaviour is shaped by a tendency to enhance or prolong feelings of pleasure and avoid or minimize feelings of pain (Kahneman, 1999). The present study focused upon *core affect*, which has been defined as “the most elementary consciously accessible affective feelings” (Russell & Feldman Barrett, 1999, p. 806) and is considered the most general affective construct when compared to emotion and mood (Ekkekakis, 2013).

It has been demonstrated that musical accompaniment has the capacity to influence the affective responses of exercisers and is frequently combined with visual stimuli such as video within modern exercise and sport training facilities (see Hallett & Lamont, 2015; Karageorghis & Priest, 2012a, 2012b; Loizou, Karageorghis, & Bishop, 2014). A distinction can be made between music-and-video and music-video in these settings. Music-and-video can be considered an umbrella term for all instances in which auditory and visual stimuli are experienced concurrently. The use of incongruent stimuli (e.g., music combined with rural scenes) has been the subject of recent investigation (e.g., Barwood, Weston, Thelwell, & Page, 2009; Jones, Karageorghis, & Ekkekakis, 2014). In contrast, the present investigation focused on music-video, a specific form of music-and-video, wherein auditory and visual stimuli are congruent (i.e., the official video of the piece of music that is being played). The paucity of research examining the effects of music-video is rather surprising given its

prominence within exercise settings and there are fewer than 10 related studies in the literature (e.g., Bigliassi et al., 2014; Hutchinson, Karageorghis, & Jones, 2015). Given that individuals are more likely to participate in exercise programs that they find to be enjoyable (Greene & Petruzzello, 2015), interventions that are targeted to improve the exercise experience are likely to facilitate exercise adherence. Accordingly, insights into such interventions could hold particular value for health and exercise practitioners.

### *The circumplex model of affect*

In order to capture affective phenomena at a global level, we conceptualize core affect as a dimensional domain, characterized by affective valence (ranging from pleasure to displeasure) and perceived activation, which can be considered as two orthogonal and bipolar dimensions (Russell, 1980). Therefore, different affective states are considered to be combinations of varying degrees of these two constituent dimensions. The circumplex model of affect (Russell, 1980) has been recommended as a conceptual framework to enable measurement of the full range of core affect responses to exercise (Ekkekakis & Petruzzello, 2002) and widely used in the field of psychomusicology (e.g., Krause, North, & Hewitt, 2015; North & Hargreaves, 1997; Ritossa & Rickard, 2004). It consists of two dimensions (affective valence and perceived activation) and thus provides four quadrants (a) high-activation pleasant affect (e.g., excitement), (b) high-activation unpleasant affect (e.g., afraid), (c) low-activation unpleasant affect (e.g., sluggish) and (d) low-activation pleasant affect (e.g., peaceful). Categorical measures of affect rely on a set of predetermined scales in order to map individual differences in affect using quantitative differences in item ratings (Ekkekakis & Petruzzello, 2002). Contrastingly, the circumplex affords individuals the freedom to accurately describe their affective experiences, over a prolonged period of time,

which allows for an almost unlimited number of individual affective profiles (Ekkekakis & Petruzzello, 2002).

#### *Dual-mode theory of affective responses to exercise*

Ekkekakis (2003, 2005) proposed the dual-mode theory to depict the relationship between exercise intensity and affective responses. An underlying premise of this theory is that exercise intensity should be defined according to a fixed metabolic marker, such as the ventilatory threshold or lactate threshold (Rose & Parfitt, 2008); such markers are associated with a host of physiological changes (e.g., increased respiration rate, accumulation of lactic acid). The dual-mode theory posits that affective responses to exercise below the ventilatory threshold/lactate threshold are primarily driven by cognitive factors and are generally positive. Affective responses to exercise proximal to the ventilatory threshold/lactate threshold vary among exercisers, with some reporting positive, and others reporting negative affect. Conversely, at higher intensities above the ventilatory threshold/lactate threshold, *interoceptive* or bodily-related cues gain salience and affective responses are generally less positive as a physiological steady state becomes difficult or even impossible to maintain (Ekkekakis, Parfitt, & Petruzzello, 2011). Furthermore, upon cessation of exercise that induces a decline in pleasure (such as exercise that exceeds the ventilatory threshold/lactate threshold), the predominant response is expected to be a robust rebound towards pleasure; a phenomenon that has been referred to recently as “the affective rebound” (Ekkekakis, 2013; Jones et al., 2014).

#### *Mechanisms underlying affective responses to exercise with music*

A number of mechanisms have been advanced with reference to the affective responses to exercise with musical accompaniment (Karageorghis, 2015; Karageorghis & Priest, 2012a,

2012b). For example, listening to music can influence attentional processing. A distinction has been made between two broad categories of attentional focus, *association* and *dissociation* (Morgan & Pollock, 1977). Association is regarded as turning focus towards task-relevant stimuli, whereas dissociation is regarded as turning focus towards task-irrelevant stimuli (Stevinson & Biddle, 1998). It has been hypothesized that dissociative strategies, such as listening to music, attenuate fatigue-related sensations by occupying the limited attentional processing capacity (e.g., Clark, Baker, & Taylor, 2015; Rejeski, 1985).

Music is also capable of evoking a broad range of feelings, among which core affect plays a particularly salient role (Juslin & Västfjäll, 2008). Moreover, many of the psychological constructs that are theorized to mediate musically-induced emotions contain a visual element (Juslin, 2013). For example, *emotional contagion* describes a process whereby individuals may “catch” the emotions of others when hearing their vocal expressions or seeing their facial expressions (Juslin & Västfjäll, 2008). The use of music-video, wherein the performing artist is usually heard *and* seen, could be particularly potent in this regard.

In addition to attentional processing and emotional engagement, Mitchell and MacDonald (2006) proposed that music can imbue the listener with a sense of control. Their research focused on the phenomenon of *audioanalgesia* (i.e., the use of music to reduce the perception of pain) in clinical settings, where there is generally little perceived control over one’s immediate surroundings. The perception of control is also important within an exercise environment and can contribute towards an individual’s overall evaluation of the experience (Rose & Parfitt, 2010).

#### *Affective responses to auditory and visual stimuli during exercise*

Karageorghis, Terry, and Lane (1999) developed a conceptual framework to predict the effects of *asynchronous* or background music use in exercise and sport settings. Their

framework underscored the Brunel Music Rating Inventory (BMRI), which has been used to assess the motivational qualities of music. The BMRI was revised to create the BMRI-2 (Karageorghis, Priest, Terry, Chatzisarantis, & Lane, 2006), an instrument that was adopted for the present study and has been shown to be valid and reliable by its originators, as well as by independent researchers (e.g., Clark, Baker, Peiris, Shoebridge, & Taylor, 2015). The BMRI-2 only includes items that pertain to the audible properties of the musical stimulus, often referred to in the literature as *intrinsic* or *congeneric* music factors and is accompanied by a series of recommendations to account for some of the *extrinsic* or *extragenetic* factors (see Karageorghis et al., 2006, p. 907).

The conceptual frame underlying the study of music in exercise and sport has also been revised recently (Karageorghis, 2015). The revisions were inspired, in part, by theoretical advancements in psychomusicology involving the reciprocal feedback model (Hargreaves, MacDonald, & Miell, 2005), as well as advancements in the domain of exercise and sport (e.g., Bishop, Karageorghis, & Loizou, 2007; Karageorghis et al., 1999; Terry & Karageorghis, 2006). In particular, the Hargreaves et al. (2005) model brought to the fore the interconnectedness of the music, the listener, and the context. Karageorghis (2015) arranged *music factors* in a hierarchy of intrinsic (e.g., tempo) and extrinsic (e.g., cultural associations) factors and presented them as antecedents. *Personal factors* (e.g., gender) and *situational factors* (e.g., mode of exercise) were proposed to moderate a number of consequences that include the regulation of affective valence and perceived activation. Moreover, the new theoretical model postulates a series of reciprocal interactions and feedback loops among the antecedents, moderators, and consequences of music use that determine future selection decisions.

Researchers have examined the impact of music on affective responses under varying situational conditions, by employing numerous exercise modalities such as treadmill running

(Terry, Karageorghis, Mecozzi Saha, & D'Auria, 2012), group-based classes (Clark, Baker, & Taylor, 2012), and cycle ergometry (Lim, Karageorghis, Romer, & Bishop, 2014).

Comparatively, there is a distinct lack of research focused on the mediating role of the listener's personal characteristics, such as gender, in the context of musically accompanied exercise (Karageorghis & Priest, 2012b). A study by Karageorghis et al. (2010) provides a notable exception given that females were found to derive greater pleasure than their male counterparts when exercising to exhaustion with musical accompaniment. However, in this instance, participants were required to synchronize their movements to the tempo of the music; a motor task that might have been more familiar to the female participants (see, e.g., Cain et al., 2015). There is a strong case for additional research to uncover any possible gender differences in the response to asynchronous music (i.e., without auditory-motor synchronization) during exercise.

Karageorghis et al. (2009) compared motivational music to a no-music control during a treadmill task and reported significantly enhanced affect during the music condition. Furthermore, Elliot, Carr, and Savage (2004) compared motivational music to *oudeterous* (neutral in motivational terms; see Karageorghis et al., 1999) music and a no-music control condition during a 12-min cycle ergometry protocol. Participants reported significantly higher positive affect during the motivational condition compared to the no-music control. However, the exercise intensity adopted in both of the aforementioned studies warrants further examination. Defining exercise intensity according to the ventilatory threshold/lactate threshold is preferable to the use of percentage of heart rate max (HRmax) or a fixed rating of perceived exertion (RPE; Elliot et al., 2004; Karageorghis et al., 2009). This approach ensures much less inter-individual variability in metabolic state when participants respond to affective measures during exercise (Rose & Parfitt, 2012), thus removing a potential



confound when comparing affective responses across individuals and between intensities (Rose & Parfitt, 2010).

Despite the suggestion that the interactive effects of music and visual stimulation are more potent in combination than when presented in isolation (Baumgartner, Lutz, Schmidt, & Jäncke, 2006; Loizou et al., 2014), research examining the interactive effects of music-and-video accompaniment to exercise is scant. The limited findings reveal that music-and-video interventions can promote greater attendance in exercisers at a health club over a 14-week period (Annesi, 2001), increase work output (Barwood et al., 2009) and elicit more positive core affect (Jones et al., 2014). There are numerous limitations associated with such studies, such as the lack of standardization in terms of music intensity (volume) and exercise duration (Annesi, 2001), as well as incongruence between auditory and visual stimuli (Jones et al., 2014). Moreover, Barwood et al.'s (2009) experiment was conducted in a climatic chamber that elicited a "moderate risk of heat illness" (p. 437) and is not representative of most contemporary exercise facilities, which offer climate control.

Recently, three groups of researchers have investigated the impact of music-video accompaniment during exercise (Bigliassi et al., 2014; Hutchinson et al., 2015; Lin & Lu, 2013). Their findings indicate that music-video can reduce participants' RPE (Bigliassi et al., 2014; Hutchinson et al., 2015; Lin & Lu, 2013), promote a dissociative attentional focus (Hutchinson et al., 2015), increase work output (Lin & Lu, 2013) and enhance core affect (Hutchinson et al., 2015). Nonetheless, a number of limitations can be identified within such studies that include a failure to familiarize participants with sensory deprivation (Lin, & Lu, 2013), a lack of consideration towards the motivational qualities of music (Bigliassi et al., 2014) and not examining the influence of auditory and visual stimuli directly at the ventilatory threshold (e.g., Hutchinson et al., 2015). There is, therefore, considerable scope to

examine the effects of music-video accompaniment to exercise while accounting for such limitations.

### *Aims and hypotheses*

To date, only two attempts have been made to examine the influence of music-and-video on core affect during and post-exercise at an intensity that is fully standardized across participants (Hutchinson et al., 2015; Jones et al., 2014). In these studies, the researchers used intensities below and above ventilatory threshold. However, to our knowledge, no study to date has investigated such effects at the lactate threshold; the exercise intensity that was adopted for the present study. Core affect is open to individual interpretation and thus relatively malleable when exercising at lactate threshold (Ekkekakis, 2003, 2013). Given the call for pleasure to be added as the “third pillar” of exercise prescription (Guiraud, Labrunee, Gayda, Juneau, & Gremeaux, 2012), interventions aimed to enhance the experience of exercise could prove vital for health and exercise practitioners in their work to combat physical inactivity. The primary aim of the present study was to examine the influence of music and music-video on affective responses during and immediately following cycle ergometry conducted at the lactate threshold. The secondary aim was to examine the degree to which gender moderated the influence of music and music-video on affective responses.

We tested four hypotheses: (a) participants’ affective valence would be most positive (situated in the right-hand quadrants of the circumplex model of affect) in the music and music-video conditions, and least in control ( $H_1$ ); (b) participants’ perceived activation would be the highest (situated in the upper quadrants of the circumplex model of affect) in the music and music-video conditions, and lowest in control ( $H_2$ ); (c) the control condition would be associated with the greatest affective rebound compared to both experimental conditions, owing to participants’ more positive in-task affective state in the music and music-video

conditions ( $H_3$ ); and (d) gender would not moderate the influence of music and music-video on affective responses either during or immediately after exercise ( $H_4$ ).

## **Materials and methods**

### *Study design*

There were three conditions that were administered using a randomized, fully counterbalanced, crossover design.

### *Participants*

Eligible participants for experimental trials were adults aged 18 years or over who were able to participate safely in exercise in accord with responses to the Physical Activity Readiness Questionnaire (PAR-Q; Canadian Society for Exercise Physiology, 2002). Participants were deemed ineligible if they had a hearing deficiency and/or any visual impairment that could not be overcome through the use of spectacles or contact lenses. Twenty-four undergraduate students were screened for eligibility and recruited (eight women and 16 men;  $M_{\text{age}} = 21.3$  years,  $SD = 1.2$  years). Participants were recreationally active, self-reporting engagement in exercise of more than 4 hr per week ( $M_{\text{training}} = 4.4$  hr per week,  $SD = 1.5$  hr per week). Each participant was provided with an information sheet and signed an informed consent form. The study took place at two universities located in the south of England, UK, the review boards of which provided ethical approval.

### *Interventions*

*Music and music-video selection.* To apply some objectivity in the selection of music for the experimental conditions, 10 volunteers (four women and six men) who were homogenous in

terms of age ( $M_{\text{age}} = 22.0$  years,  $SD = 1.0$  years) supplied three tracks that they considered to be “motivational” for the purpose of stationary cycling. One of the experimenters assessed the motivational qualities of the music tracks using the BMRI-2 (Karageorghis et al., 2006) to ensure that the music used in experimental trials was relatively homogenous in terms of its motivational qualities. The six tracks with the highest motivational quotients were selected: *Levels* by Avicci (126 bpm), *Titanium* by David Guetta ft. Sia (126 bpm), *Call On Me* by Eric Prydz (126 bpm), *Sweet Nothing* by Calvin Harris ft. Florence Welch (128 bpm), *Human* by The Killers (136 bpm), and *Zombie Nation* by Kernkraft 400 (140 bpm). The official music-videos of the aforementioned tracks were selected for use in the experimental trials.

*Preliminary testing.* The lactate threshold is the “lowest workload at which the rate of lactate appearance in the blood starts to exceed the rate of removal” (Ekkekakis, Hall, & Petruzzello, 2005, p. 482). Each participant was administered an incremental cycle ergometry test, in which the load was increased by .5 kg every 4 min and blood samples were taken from the fingertip during the final min of each stage in order to calculate lactate threshold. Blood samples were analyzed immediately. The D-max method (Cheng et al., 1992) was used to determine lactate threshold, which has high test-retest reliability and correlation coefficients of .77–.93 (Zhou & Weston, 1997). The D-max was taken as the point at which the slope of the polynomial curve was equal to the slope of the straight line and the appropriate load for the experimental trials was calculated from the lactate threshold. Participants’ affective responses were measured every 4 min during preliminary testing to aid familiarization.

*Experimental trial.* Experimental testing took place at least 48 hr after the preliminary testing. Participants visited the laboratory on three separate occasions to be exposed to three conditions (a) music, (b) music-video, and (c) control (no music or music-video).

Experimental trials took place a minimum of 48 hr apart. Participants were required to avoid caffeine and alcohol, maintain adequate hydration, and gain sufficient sleep during the night prior to testing. Upon entry to the laboratory, each participant had a 5-min period of seated rest, a 5-min warm-up on a cycle ergometer, and then a 20-min cycle at their lactate threshold. Each participant completed a 5-min active cool-down on the cycle ergometer that was followed by a 15-min period of seated rest.

*Apparatus.* A cycle ergometer (Monark 874E) was used for testing and blood samples (to determine lactate threshold) were analyzed using a lactate analyzer (Biosen Line-C). Music was administered via over-ear headphones (Sony MDR-V150) at an approximate intensity of 70 dBA, which is deemed safe from an audiological perspective (Health and Safety Executive, 2005). Music and music-video were delivered via a media player (VLC 2.1.0) on a laptop (Toshiba Satellite Pro C850-15T; screen size 15.6 in), placed 1.5 m in front of the participant, slightly below eye level. The headphones and laptop were not used when the control condition was administered in order to maintain external validity.

### *Outcomes*

*Core affect measurement.* On the basis that we conceptualize affect as a dimensional domain, affective valance was assessed using Hardy and Rejeski's (1989) Feeling Scale, which has an 11-point scale ranging from (I feel) *very good* (+5) to *very bad* (-5). This is an exercise-specific measure that has had its validity demonstrated in a series of studies (Hardy & Rejeski, 1989) and been used extensively in exercise-related research (e.g., Rose & Parfitt, 2008). Perceived activation was measured using the Felt Arousal Scale (FAS; Svebak & Murgatroyd, 1985), which is a single-item, 6-point measure ranging from 1 (*low arousal*) to 6

(*high arousal*). The validity of the FAS has been demonstrated by van Landuyt, Ekkekakis, Hall, and Petruzzello (2000) and it has been used extensively in the exercise context (e.g., Rose & Parfitt, 2012). Baseline affect was measured following the 5-min period of seated rest. In-task core affect was measured every 2 min during the exercise. Post-task core affect was measured at 0, 5, 10, and 15 min following completion of the cool-down.

### *Sample size*

A power analysis was conducted using G\*Power 3 (Faul, Erdfelder, Buchner, & Lang, 2009) to determine an appropriate sample size. Based on a large predicted effect size derived from Hutchinson et al. (2015;  $\eta_p^2 = .32$ ), an alpha level of .05, and power at .85, the analysis indicated that 23 participants would be required. To minimize the impact of possible participant attrition and multivariate outliers, one additional participant was recruited.

### *Data analysis*

The data were screened for univariate and multivariate outliers and relevant checks were undertaken to ensure that the data met the parametric assumptions that underlie mixed-model MANOVA. Furthermore, a variety of data transformations were planned for (e.g., square root, logarithm) in case that any of the data needed to be normalized. Pre-task affective measures were examined and planned to be used as covariates if statistical differences emerged across conditions. The affective variables (affective valence and perceived activation) were assessed using a 3 (condition) x 5 (time) x 2 (gender) MANOVA. The means of three in-task time-point ranges were selected for statistical analysis: 2–6 min, 8–14 min and 16–20 min. Post-task time-points selected for statistical analysis were at 0 and 5 min post cool-down. To facilitate a visual interpretation of the findings, the FS and FAS scores were plotted using the circumplex model of affect (Russell, 1980).

## Results

There were no outliers identified and the data met most of the assumptions that underlie mixed-model MANOVA. Tests of the distributional properties of the data in each cell of the analysis revealed violations of normality in 13 of the 60 cells (five at  $p < .05$ , three at  $p < .01$ , and five at  $p < .001$ ). The sphericity assumption was violated in all  $F$  tests and Greenhouse-Geisser adjustments were made accordingly.

### *Interaction effects*

There was no significant higher-order interaction for affective valence or perceived activation (condition x time x gender; see Table 1). There was, however, a two-way interaction for affective valence ( $p < .001$ ,  $\eta_p^2 = .20$ ; condition x time; see Table 1), associated with a large effect size. The music and music-video conditions elicited higher affective valence than the control condition at the first two time points during the exercise bout (see Figure 1a).

Furthermore, the music-video condition elicited higher affective valence than the music condition at these time-points. There were no significant differences in affective valence for the final in-task measurement point. Immediately following the cool-down, affective valence was significantly higher in the music and music-video conditions when compared to control.

There were no significant differences in affective valence across conditions at 5 min post-task (see Figure 1a). There was a two-way interaction effect for perceived activation ( $p = .020$ ,  $\eta_p^2 = .17$ ; gender x time; see Table 1), associated with a large effect size, wherein perceived activation was higher for males than females at the final in-task measurement point, and significantly lower for males than females at cool-down and 5 min post-task.

**\*Insert Table 1 about here\***

**\*Insert Figure 1 about here\***

*Main effects*

There was a main effect of condition for affective valence ( $p < .001$ ,  $\eta_p^2 = .40$ ; see Table 1), associated with a large effect size. Pairwise comparisons showed that affective valence was significantly higher for music when compared to control ( $p = .001$ , 95% CI [0.47, 1.94]) and for music-video when compared to control ( $p = .002$ , 95% CI [0.47, 2.21]). There was a main effect of time for both affective valence ( $p < .001$ ,  $\eta_p^2 = .48$ ) and perceived activation ( $p < .001$ ,  $\eta_p^2 = .62$ ; see Table 1), both associated with a large effect size. Affective valence scores decreased incrementally throughout the exercise bout, and then increased immediately after the exercise bout. Affective valence scores were significantly higher 5 min post-task compared to the start of the exercise bout ( $p = .005$ , 95% CI [0.30, 2.21]). Perceived activation scores increased incrementally throughout the exercise bout and then decreased incrementally following the exercise bout. Perceived activation scores were significantly lower 5 min post-task than at the start of the exercise bout ( $p < .001$ , 95% CI [-1.20, -0.30]).

**Discussion**

The main purpose of this study was to examine the influence of music and music-video on core affect (affective valence and perceived activation) during and immediately following cycle ergometry at the lactate threshold. A secondary purpose was to investigate the degree to which gender moderated the influence of music and music-video on affective responses.

Results indicated that there was a main effect of condition for affective valence (see Table 1).

The music and music-video conditions facilitated more positive affective responses than the control condition, thus  $H_1$  was accepted. The hypothesis pertaining to perceived activation ( $H_2$ ) was not accepted given that there was no main effect of condition (see Table 1).

Following the exercise bout, the affective rebound was largest in the control condition and therefore  $H_3$  was accepted. Finally,  $H_4$  was accepted given that gender did not moderate the



influence of music and music-video on affective responses either during or immediately after exercise. The present findings should be interpreted with caution given the violations of normality that were evident in the data (13 out of 60 cells [21.7%]).

### *Affective variables*

*Affective valence.* The present findings indicated that motivational music can elicit more positive affective valence when compared to a control condition, which is echoed in previous research (Karageorghis et al., 2009; Lim et al., 2014; Terry et al., 2012). Moreover, the findings illustrate how the addition of a visual stimulus further enhanced participants' affective responses during exercise at the lactate threshold, an intensity at which affective responses are hypothesized to be the most malleable (Ekkekakis et al., 2011).

When exercising at, or close to the lactate threshold, internal and external cues compete, given that the capacity for attentional processing is reduced (Tenenbaum, 2001). It is plausible that the music-video condition provided a more salient external cue and consequently promoted a more dissociative attentional focus when compared with music-only. Thus, the fatigue-related sensations associated with exercising at the lactate threshold could have been less prevalent in the music-video condition, resulting in a higher affective valence score (cf. Hutchinson & Karageorghis, 2013). What is noteworthy upon a closer examination of the condition x time interaction is the convergence in affective scores at the end of the exercise bout in the two experimental conditions. There is a larger difference evident, albeit non-significant, between the music-only and music-video conditions at the first two in-task measurement points (2–6 min;  $M_{diff} = .38$ ; 8–14 min;  $M_{diff} = .51$ ) when compared to the final in-task measurement point (16–20 min;  $M_{diff} = .18$ ). This is indicative of the reduced attentional processing capacity that is available to exercisers during the final

stages of exercise (Tenenbaum, 2001). The benefits reported herein with regard to enhanced affect, are amplified by other recent studies that have investigated the impact of music-video on a range of performance-related and psychophysical variables (Bigliassi et al., 2014; Hutchinson et al., 2015; Lin & Lu, 2013). Accordingly, it is evident that music-video has the potential to influence a range of outcomes that are relevant both to physical performance and health maintenance (cf. Clark et al., 2015; Hallett & Lamont, 2015).

The present findings illustrate the potential efficacy of music-video in enhancing core affect; a benefit of music-only use that has been known about in the exercise domain for some time (see, e.g., Karageorghis & Terry, 1997). Given that the tracks were performed or visually depicted in the music-videos, their emotional salience was heightened and there is thus greater scope for the phenomenon of emotional contagion to occur (Juslin & Västfjäll, 2008). Notably, the findings indicate that, post-task, music elicits more positive core affect than music-video ( $M_{diff} = .28$ ). The cessation of exercise was followed by an improvement in affective valence during the cool-down phase and 5 min post-exercise in all conditions (see Figure 1a). Nonetheless, the control condition was associated with the most pronounced improvement; affective valence scores were elevated towards those observed in the experimental conditions. This has been referred to as the “rebound phenomenon” and repeatedly identified in the affect literature (e.g., Greene & Petruzzello, 2015; Stych & Parfitt, 2011). The finding is consistent with the dual-mode theory (Ekkekakis, 2003, 2005), which postulates that a robust rebound towards pleasure is expected following the termination of exercise that has induced a decline in affect.

*Perceived activation.* The experimental conditions had little bearing on perceived activation and a close examination of the findings indicates that participants reported the highest perceived activation in the music condition (see Figure 1b). This finding is consistent with

recent research that has examined exercise intensities above the ventilatory threshold (Hutchinson et al., 2015; Jones et al., 2014). Many researchers in this field have suggested that the most-frequently cited goal of music is to influence feelings (Juslin & Laukka, 2004; Laukka & Quick, 2013). In a review of the psychological constructs that mediate musically-induced feelings, Juslin (2013) explained that *visual imagery* refers to a process whereby feelings are evoked in the listener as a result of the inner images that they conjure up while listening to music. Thus, it is plausible that the physical images contained within the music-videos were deemed to be less stimulating than the participants' internal images when exercising in the music and control conditions.

The significant main effect of time revealed that all conditions elicited a decrease in perceived activation from the latter stages of the exercise bout to post 5-min (see Figure 1b). This is in accordance with the average post-exercise state that is characterized by a gradual decrease in perceived activation (Ekkekakis, 2003). It also points to a specialist role that music might play in expediting post-exercise recovery through the principle of *entrainment*, wherein the main pulses of the body such as respiration and heart rate can be “guided” towards resting levels by appropriate musical stimuli (see Karageorghis, 2015).

*Moderating influence of gender.* The non-significant higher-order interaction for affective valence and perceived activation revealed that gender did not moderate the influence of music and music-video on affective responses during and immediately post-exercise. The present results are thus dissimilar to those of Karageorghis et al. (2010) who reported that women exhibited significantly higher affective valence scores than men when exposed to music. The discrepancy in findings might be attributed to differing exercise modalities (cycle ergometry vs. circuit-type exercises to exhaustion), as well as the manner in which the music was used (asynchronous vs. synchronous application). Both of these are task-related

situational factors that were theorized to moderate affective responses to music in a recent model (Karageorghis, 2015). For perceived activation, there was a significant ( $p = .020$ ) gender x time interaction wherein FAS scores significantly increased during and decreased post-exercise for men only. One might speculate that this difference has an evolutionary antecedent: Males generally experience stronger psychobiological responses to physical activity given the salience of such responses to the hunting and defending conducted by our male forebears (Lombardo, 2012).

#### *Limitations of the present study*

As previously highlighted, there were multiple violations of normality evident in the affective data. Planned transformations would only have served to normalize a proportion of the non-normal data, while rendering other sections of the dataset non-normal. Accordingly, we decided not to run transformations, and the results of the present study should be interpreted with these violations in mind.

We compared affective responses to music and music-video during exercise at the lactate threshold. A further limitation was that a video-only condition was not included, therefore the capacity of video alone in mediating affective responses during- and post-exercise remains unexplored. Nonetheless, in naturalistic conditions such as health and fitness facilities, it is unlikely that exercisers would watch a music-video without the accompanying sound (see also Hutchinson et al., 2015). Moreover, we did not measure participants' responses to music and music-video while they were at rest. This was due to a previous suggestion that engagement in exercise influences how individuals respond to music (Karageorghis & Jones, 2014). However, in exercise and health facilities it is highly likely that such stimuli would be present both prior to and following an exercise bout.

Another limitation of the present study was that the music used in the experimental trials was subject to specific selection criteria but the same cannot be said for the video content. Ten volunteers each provided three tracks that they considered to be motivational for stationary cycling and the tracks were subsequently rated by one of the experimenters using the BMRI-2 (Karageorghis et al., 2006). The experimental participants were not afforded the opportunity to rate the music themselves and thus the extent to which *they* found it motivating for cycle ergometry remains unknown. It has been argued that experimenter-selected music can compromise the perception of control for the listener, which has been identified as one of the mechanisms underlying the efficacy of music in regulating affective responses (Chanda & Levitin, 2013; Mitchell & MacDonald, 2006).

The study was conducted in a laboratory environment and therefore the results may not be directly transferable to real-world settings (Glasgow, Green, & Ammerman, 2007). Indeed, cognitive factors may influence affective responses when exercising in stimulating environments (e.g., an exercise facility or outdoors) when compared to laboratory settings (Rose & Parfitt, 2010). Generalization of the present findings to the wider population may also be limited by the homogenous characteristics of the sample employed (i.e., age and training status), as well as the exercise modality (cycle ergometry).

### *Practical implications*

The present findings provide empirical evidence to support the efficacy of music and music-video as a strategy for enhancing the exercise experience. This is illustrated in Figure 1c wherein only the control condition passed through the low-activation unpleasant quadrant of the circumplex model of affect (Russell, 1980). Affective responses to exercise are generally positive at low intensities, without the need for affect-enhancing strategies (Ekkekakis et al., 2011). However, the potential health benefits (e.g., cardio-respiratory fitness) associated with

exercise are more substantial when higher-intensity exercise is performed (Riddoch & O'Donovan, 2006), albeit that this is subject to greater inter-individual variability in affect (Ekkekakis, 2003). Researchers have demonstrated the role of affective responses (both during and after exercise) in facilitating adherence to exercise (Williams, Dunsiger, Jennings, & Marcus, 2012). Habitual engagement in exercise confers several benefits in terms of physical health and mental wellbeing (Biddle et al., 2010). Likewise, at a societal level, raising adherence to exercise is advantageous given the considerable cost burden that has been associated with sedentary lifestyles (Townsend, Wickramasinghe, Williams, Bhatnagar, & Rayner, 2015). The key message for exercise or music therapy practitioners is that music and music-video may contribute to enhanced public health by increasing the appeal of higher-intensity exercise and facilitating adherence.

The affective rebound experienced post-exercise by those in the music condition is also noteworthy (see Figure 1a). Practitioners are urged to consider the use of music in parts of exercise facilities that are used outside of the physical workout (e.g., changing facilities, onsite cafes) to enhance the overall experience of exercise. An added benefit to employing both music and music-video within exercise settings is its relative ease of implementation (Jones et al., 2014). From a practical perspective, instructor-led exercise programmes could be delivered via video and combined with music in order to promote positive affective responses to exercise. This is an inexpensive means by which to deliver such programmes, and would allow individuals to exercise at a time and place that is convenient to them. Moreover, there is scope for music and video choices to be made with direct reference to the circumplex model of affect in order that a *desired* affective state can be engendered by the combination of auditory and visual stimuli (cf. North & Hargreaves, 1997; Ritossa & Rickard, 2004).

## Conclusions and recommendations

This is the first study to demonstrate that music-videos can enhance core affect when exercising at the lactate threshold. From a theoretical standpoint, these findings are important because exercise at intensities proximal to the lactate threshold is associated with considerable affective variability (Ekkekakis, 2003, 2013). A strength of this study is that it focused exclusively on affective responses to exercise. Conversely, it has been suggested that the invasive nature of examining physiological measures (e.g., blood lactate levels) has the potential to negatively influence the effects of music (Karageorghis & Priest, 2012a).

A useful addition to this line of research would be the development of a method for selecting the most appropriate music-videos to accompany exercise. Future researchers may also consider incorporating a video-only and/or music-video-primed condition (see, e.g., Loizou & Karageorghis, 2015) so that a more comprehensive comparison can be made between auditory and visual stimuli. Moreover, it would be worthwhile to assess the influence of such stimuli on affective responses before, during, and post-exercise, given the likelihood of such exposure within contemporary exercise facilities. Additional research is necessary to explore and explain the varying influences of music-video between the genders. Affective responses differ according to individuals' previous exposure to exercise (Stych & Parfitt, 2011), therefore future researchers might also consider replicating the present study with a previously inactive sample.

In conclusion, it appears that music and music-video can be used as a valuable tool to enhance affective responses to exercise. Given the emerging empirical support pertaining to a positive relationship between affective responses to exercise and adherence (e.g., Williams et al., 2012), consideration of the music and music-videos played within exercise facilities is warranted and has positive implications for health and exercise practitioners. It is not only what one *hears* during exercise but also what one *sees* that can have a potent influence upon how they feel.

## References

- Annesi, J. J. (2001). Effects of music, television, and a combination entertainment system on distraction, exercise adherence, and physical output in adults. *Canadian Journal of Behavioural Science, 33*, 193–201.
- Barwood, M. J., Weston, J. V., Thelwell, R., & Page, J. (2009). A motivational music and video intervention improves high-intensity exercise performance. *Journal of Sports Science & Medicine, 8*, 435–442.
- Baumgartner, T., Lutz, K., Schmidt, C. F., & Jäncke, L. (2006). The emotional power of music: How music enhances the feeling of affective pictures. *Brain Research, 1075*, 151–164.
- Biddle, S. J. H., Cavill, N., Ekelund, U., Gorely, T., Griffiths, M., Jago, R., ... Richardson, D. (2010). *Sedentary behaviour and obesity: Review of the current scientific evidence*. London, UK: Department of Health. Retrieved from [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/213745/dh\\_128225.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213745/dh_128225.pdf)
- Bigliassi, M., Storti Peruzollo, A., Ferreira Dias Kanthack, T., Barreto da Silva, V., Pezarat-Correia, P., & Atimari, L. R. (2014). Effects of a sensory strategy in an isometric muscular endurance task. *Revista Andaluza de Medicina del Deporte, 7*, 55–59.
- Bishop, D. T., Karageorghis, C. I., & Loizou, G. (2007). A grounded theory of young tennis players' use of music to manipulate emotional state. *Journal of Sport & Exercise Psychology, 29*, 584–607.
- Cain, K. L., Gavand, K. A., Conway, T. L., Peck, E., Bracy, N. L., Bonilla, E., ... Sallis, J. F. (2015). Physical activity in youth dance classes. *Pediatrics, 135*, 1066–1073.
- Canadian Society for Exercise Physiology. (2002). Physical Activity Readiness Questionnaire (PAR-Q). Ontario, CAN: CSEP. Retrieved from <https://www.csep.ca/CMFiles/publications/parq/par-q.pdf>



- Chanda, M. L., & Levitin, D. J. (2013). The neurochemistry of music: Evidence for health outcomes. *Trends in Cognitive Sciences, 17*, 179–193.
- Cheng, B., Kuipers, H., Snyder, A. C., Keizer, H. A., Jeukendrup, A., & Hesselink, M. (1992). A new approach for the determination of ventilatory and lactate thresholds. *International Journal of Sports Medicine, 13*, 518–522.
- Chief Medical Officers of England, Scotland, Wales and Northern Ireland. (2011). *Start active, stay active: A report on physical activity from the four home countries' Chief Medical Officers*. London, UK: Department of Health. Retrieved from [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/216370/dh\\_128210.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216370/dh_128210.pdf)
- Clark, I. N., Baker, F. A., & Taylor, N. F. (2012). The effects of live patterned sensory enhancement on group exercise participation and mood in older adults in rehabilitation. *Journal of Music Therapy, 49*, 180–204.
- Clark, I. N., Baker, F. A., & Taylor, N. F. (2015). The modulating effects of music listening on health-related exercise and physical activity in adults: A systematic review and narrative synthesis. *Nordic Journal of Music Therapy*. Advance online publication. doi:10.1080/08098131.2015.1008558
- Clark, I. N., Baker, F. A., Peiris, C. L., Shoebridge, G., & Taylor, N. F. (2015). The Brunel Music Rating Inventory-2 is a reliable and valid instrument for older cardiac rehabilitation patients selecting music for exercise. *Psychology of Music*. Advance online publication. doi:10.1177/0305735614565830
- Ekkekakis, P. (2003). Pleasure and displeasure from the body: Perspectives from exercise. *Cognitive and Emotion, 17*, 213–239.

- Ekkekakis, P. (2005). The study of affective responses to acute exercise: The dual-mode model. In R. Stelter & K. K. Roessler (Eds.), *New approaches to sport and exercise psychology* (pp. 119–146). Oxford, UK: Meyer & Meyer Sport.
- Ekkekakis, P. (2013). Pleasure from the exercising body: Two centuries of changing outlooks in psychological thought. In P. Ekkekakis (Ed.), *Routledge handbook of physical activity and mental health* (pp. 35–56). New York, NY: Routledge.
- Ekkekakis, P., & Dafermos, M. (2012). Exercise is a many-splendored thing, but for some it does not feel so splendid: Staging a resurgence of hedonistic ideas in the quest to understand exercise behavior. In E. O. Acevedo (Ed.), *Oxford handbook of exercise psychology* (pp. 295–333). New York, NY: Oxford University Press.
- Ekkekakis, P., Hall, E. E., & Petruzzello, S. J. (2005). Variation and homogeneity in affective responses to physical activity of varying intensities: An alternative perspective on dose-response based on evolutionary considerations. *Journal of Sport Sciences, 23*, 477–500.
- Ekkekakis, P., Parfitt, G., & Petruzzello, S. J. (2011). The pleasure and displeasure people feel when they exercise at different intensities: Decennial update and progress towards a tripartite rationale for exercise intensity prescription. *Sports Medicine, 41*, 641–671.
- Ekkekakis, P., & Petruzzello, S. J. (2002). Analysis of the affect measurement conundrum in exercise psychology: IV. A conceptual case for the affect circumplex. *Psychology of Sport and Exercise, 3*, 35–63.
- Elliott, D., Carr, S., & Savage, D. (2004). Effects of motivational music on work output and affective responses during sub-maximal cycling of a standardized perceived intensity. *Journal of Sport Behavior, 27*, 134–147.

- Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using G\*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods, 41*, 1149–1160.
- Glasgow, R. E., Green, L. W., & Ammerman, A. (2007). A focus on external validity. *Evaluation & the Health Professions, 30*, 115–117.
- Greene, D. R., & Petruzzello, S. J. (2015). More isn't necessarily better: Examining the intensity–affect–enjoyment relationship in the context of resistance exercise. *Sport, Exercise, and Performance Psychology, 4*, 75–87.
- Guiraud, T., Labrunee, M., Gayda, M., Juneau, M., & Gremeaux, V. (2012). Non-pharmacological strategies in cardiovascular prevention: 2011 highlights. *Annals of Physical and Rehabilitation Medicine, 55*, 342–374.
- Hallett, R., & Lamont, A. (2015). How do gym members engage with music during exercise? *Qualitative Research in Sport, Exercise and Health, 7*, 411–427.
- Hardy, C. J., & Rejeski, W. J., (1989). Not what, but how one feels: The measurement of affect during exercise. *Journal of Sport & Exercise Psychology, 11*, 304–317.
- Hargreaves, D. J., MacDonald, R. A. R., & Miell, D. E. (2005). How do people communicate using music? In D. E. Miell, R. A. R. MacDonald, & D. J. Hargreaves (Eds.), *Musical communication* (pp. 1–25). Oxford, UK: Oxford University Press.
- Health and Safety Executive (2005). *The Control of Noise at Work Regulations 2005*. Retrieved from <http://www.legislation.gov.uk/ukxi/2005/1643/contents/made>
- Hutchinson, J. C., & Karageorghis, C. I. (2013). Moderating influence of dominant attentional style and exercise intensity on responses to asynchronous music. *Journal of Sport & Exercise Psychology, 35*, 625–643.

- Hutchinson, J. C., Karageorghis, C. I., & Jones, L. (2015). See hear: Psychological effects of music and music-video during treadmill running. *Annals of Behavioral Medicine, 49*, 199–211.
- Jones, L., Karageorghis, C. I., & Ekkekakis, P. (2014). Can high-intensity exercise be more pleasant? Attentional dissociation using music and video. *Journal of Sport & Exercise Psychology, 36*, 528–541.
- Juslin, P. N. (2013). From everyday emotions to aesthetic emotions: Towards a unified theory of musical emotions. *Physics of Life Reviews, 10*, 235–266.
- Juslin, P. N., & Laukka, P. (2004). Expression, perception, and induction of musical emotions: A review and a questionnaire study of everyday listening. *Journal of New Music Research, 33*, 217–238.
- Juslin, P. N., & Västfjäll, D. (2008). Emotional responses to music: The need to consider underlying mechanisms. *Behavioral and Brain Sciences, 31*, 559–621.
- Kahneman, D. (1999). Objective happiness. In D. Kahneman, E. Diener, & N. Schwarz (Eds.), *Well-being: Foundations of hedonic psychology* (pp. 3–25). New York, NY: Russell-Sage.
- Karageorghis, C. I. (2015). The scientific application of music in exercise and sport: Towards a new theoretical model. In A. M. Lane (Ed.), *Sport and exercise psychology* (2nd ed., pp. 276–322). London, UK: Routledge.
- Karageorghis, C. I., & Jones, L. (2014). On the stability and relevance of the exercise heart rate–music-tempo preference relationship. *Psychology of Sport and Exercise, 15*, 299–310.
- Karageorghis, C. I., Mouzourides, D. A., Priest, D. L., Sasso, T. A., Morrish, D. J., & Walley, C. L. (2009). Psychophysical and ergogenic effects of synchronous music during treadmill walking. *Journal of Sport & Exercise Psychology, 31*, 18–36.

- Karageorghis, C. I., & Priest, D. L. (2012a). Music in the exercise domain: A review and synthesis (Part I). *International Review of Sport and Exercise Psychology*, 5, 44–66.
- Karageorghis, C. I., & Priest, D. L. (2012b). Music in the exercise domain: A review and synthesis (Part II). *International Review of Sport and Exercise Psychology*, 5, 67–84.
- Karageorghis, C. I., Priest, D. L., Terry, P. C., Chatzisarantis, N. L. D., & Lane, A. M. (2006). Redesign and initial validation of an instrument to assess the motivational qualities of music in exercise: The Brunel Music Rating Inventory-2. *Journal of Sports Sciences*, 24, 899–909.
- Karageorghis, C. I., Priest, D. L., Williams, L. S., Hirani, R. M., Lannon, K. M., & Bates, B. J. (2010). Ergogenic and psychological effects of synchronous music during circuit-type exercise. *Psychology of Sport and Exercise*, 11, 551–559.
- Karageorghis, C. I., & Terry, P. C. (1997). The psychophysical effects of music in sport and exercise: A review. *Journal of Sport Behavior*, 20, 54–68.
- Karageorghis, C. I., Terry, P. C., & Lane, A. M. (1999). Development and initial validation of an instrument to assess the motivational qualities of music in exercise and sport: The Brunel Music Rating Inventory. *Journal of Sports Sciences*, 17, 713–724.
- Krause, A. E., North, A. C., & Hewitt, L. Y. (2015). Music-listening in everyday life: Devices and choice. *Psychology of Music*, 43, 155–170.
- Laukka, P., & Quick, L. (2013). Emotional and motivational uses of music in sports and exercise: A questionnaire study among athletes. *Psychology of Music*, 41, 198–215.
- Lim, H. B. T., Karageorghis, C. I., Romer, L. M., & Bishop, D. T. (2014). Psychophysiological effects of synchronous versus asynchronous music during cycling. *Medicine & Science in Sports & Exercise*, 46, 407–413.

- Lin, J. H., & Lu, F. J. H. (2013). Interactive effects of visual and auditory intervention on physical performance and perceived effort. *Journal of Sports Science & Medicine, 12*, 388–393.
- Loizou, G., & Karageorghis, C. I. (2015). Effects of psychological priming, video, and music on anaerobic exercise performance. *Scandinavian Journal of Medicine & Science in Sports, 25*, 909–920.
- Loizou, G., Karageorghis, C. I., & Bishop, D. T. (2014). Interactive effects of video, priming, and music on emotions and the needs underlying intrinsic motivation. *Psychology of Sport and Exercise, 15*, 611–619.
- Lombardo, M. P. (2012). On the evolution of sport. *Evolutionary Psychology, 10*, 1–28.
- Mitchell, L. A., & MacDonald, R. A. R. (2006). An experimental investigation of the effects of preferred and relaxing music listening on pain perception. *Journal of Music Therapy, 43*, 295–316.
- Morgan, W. P., & Pollock, M. L. (1977). Psychologic characterization of the elite distance runner. *Annals of the New York Academy of Sciences, 301*, 382–403.
- North, A. C., & Hargreaves, D. J. (1997). Liking, arousal potential, and the emotions expressed by music. *Scandinavian Journal of Psychology, 38*, 45–53.
- Rejeski, W. J. (1985). Perceived exertion: An active or passive process? *Journal of Sport Psychology, 7*, 371–378.
- Riddoch, C., & O'Donovan, G. (2006). Moderate intensity physical activity versus vigorous activity: Today's best buy in public health. *The Sport and Exercise Scientist, 10*, 16–17.
- Ritossa, D. A., & Rickard, N. S. (2004). The relative utility of 'pleasantness' and 'liking' dimensions in predicting the emotions expressed by music. *Psychology of Music, 32*, 5–22.

- Rose, E. A., & Parfitt, G. (2008). Can the Feeling Scale be used to regulate exercise intensity? *Medicine & Science in Sports & Exercise*, *40*, 1852–1860.
- Rose, E. A., & Parfitt, G. (2010). Pleasant for some and unpleasant for others: A protocol analysis of the cognitive factors that influence affective responses to exercise. *International Journal of Behavioral Nutrition & Physical Activity*, *7*.
- Rose, E. A., & Parfitt, G. (2012). Exercise experience influences affective and motivational outcomes of prescribed and self-selected intensity exercise. *Scandinavian Journal of Medicine & Science in Sports*, *22*, 265–277.
- Russell, J. A. (1980). A circumplex model of affect. *Journal of Personality and Social Psychology*, *39*, 1161–1178.
- Russell, J. A., & Feldman Barrett, L. (1999). Core affect, prototypical emotional episodes, and other things called emotion: Dissecting the elephant. *Journal of Personality and Social Psychology*, *76*, 805–819.
- Stevinson, C. D., & Biddle, S. J. H. (1998). Cognitive orientations in marathon running and “hitting the wall”. *British Journal of Sports Medicine*, *32*, 229–235.
- Stych, K., & Parfitt, G. (2011). Exploring affective responses to different exercise intensities in low-active young adolescents. *Journal of Sport & Exercise Psychology*, *33*, 548–568.
- Svebak, S., & Murgatroyd, S. (1985). Metamotivational dominance: A multimethod validation of reversal theory constructs. *Journal of Personality and Social Psychology*, *48*, 107–116.
- Tempest, G., & Parfitt, G. (2013). Imagery use and affective responses during exercise: An examination of cerebral hemodynamics using near-infrared spectroscopy. *Journal of Sport & Exercise Psychology*, *35*, 503–513.


- Tenenbaum, G. (2001). A social-cognitive perspective of perceived exertion and exertion tolerance. In R. N. Singer, H. A. Hausenblas, & C. Janelle (Eds.), *Handbook of sport psychology* (2nd ed., pp. 810–822). New York, NY: Wiley.
- Terry, P. C. & Karageorghis, C. I. (2006). Psychophysical effects of music in sport and exercise: An update on theory, research and application. In M. Katsikitis (Ed.), *Proceedings of the 2006 Joint Conference of the Australian Psychological Society and the New Zealand Psychological Society* (pp. 415–419). Melbourne, VIC: Australian Psychological Society.
- Terry, P. C., Karageorghis, C. I., Mecozi Saha, A., & D’Auria, S. (2012). Effects of synchronous music on treadmill running among elite triathletes. *Journal of Science and Medicine in Sport, 15*, 52–57.
- Townsend, N., Wickramasinghe, K., Williams, J., Bhatnagar, P., & Rayner, M. (2015). *Physical activity statistics 2015*. London, UK: British Heart Foundation. Retrieved from <https://www.bhf.org.uk/publications/statistics/physical-activity-statistics-2015>
- Van Landuyt, L. M., Ekkekakis, P., Hall, E. E., & Petruzzello, S. J. (2000). Throwing the mountains into the lakes: On the perils of nomothetic conceptions of the exercise–affect relationship. *Journal of Sport & Exercise Psychology, 22*, 208–234.
- Williams, D. M. (2008). Exercise, affect, and adherence: An integrated model and a case for self-paced exercise. *Journal of Sport & Exercise Psychology, 30*, 471–496.
- Williams, D. M., Dunsiger, S., Jennings, E. G., & Marcus, B. H. (2012). Does affective valence during and immediately following a 10-min walk predict concurrent and future physical activity? *Annals of Behavioral Medicine, 44*, 43–51.
- Zhou, S., & Weston, S. B. (1997). Reliability of using the D-max method to define physiological responses to incremental exercise testing. *Physiological Measurement, 18*, 145–154.



**Table 1.** Descriptive Statistics and Mixed-model MANOVA for Core Affect.

Dependent Measure	Condition	2–6 During		8–14 During		16–20 During		Cool Down		Post 5	
		<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Affective Valence	Music	2.60	1.22	.75	2.59	-.60	3.75	2.19	2.02	3.78	1.06
	Music-video	2.98	.96	1.26	2.55	-.42	3.70	2.06	1.51	3.50	1.23
	Control	1.06	1.85	-.77	3.09	-1.70	3.70	.97	2.37	3.13	1.39
Perceived Activation	Music	2.72	1.01	3.51	.81	4.39	.97	2.94	.78	2.00	1.02
	Music-video	2.62	1.08	3.48	.98	4.14	1.25	2.63	.98	1.84	.99
	Control	2.69	1.00	3.53	1.13	4.16	1.52	2.78	1.01	1.94	1.09
Condition x Time x Gender interaction effects:											
FS		$F(8, 176) = 1.77, p = .128, \eta_p^2 = .07$									
FAS		$F(8, 176) = .75, p = .581, \eta_p^2 = .03$									
Condition x Time interaction effects:											
FS		$F(8, 176) = 5.40, p < .001, \eta_p^2 = .20$									
FAS		$F(8, 176) = .36, p = .867, \eta_p^2 = .02$									
Condition x Gender interaction effects:											
FS		$F(2, 44) = 3.34, p = .065, \eta_p^2 = .13$									
FAS		$F(2, 44) = 2.86, p = .082, \eta_p^2 = .12$									
Gender x Time interaction effects:											
FS		$F(4, 88) = .98, p = .359, \eta_p^2 = .04$									
FAS		$F(4, 88) = 4.50, p = .020, \eta_p^2 = .17$									
Condition main effects:											
FS		$F(2, 44) = 14.88, p < .001, \eta_p^2 = .40$									
FAS		$F(2, 44) = .55, p = .539, \eta_p^2 = .03$									
Time main effects:											
FS		$F(4, 88) = 20.48, p < .001, \eta_p^2 = .48$									
FAS		$F(4, 88) = 35.94, p < .001, \eta_p^2 = .62$									

*Note.* The *F* test for all effects for both affective valence and perceived activation was subject to Greenhouse-Geisser adjustment.  $\eta_p^2$  = partial eta squared.



**Figure 1.** Feeling Scale (FS; a) and Felt Arousal Scale (FAS; b) responses (M and SE) during all conditions, and as an affective circumplex (c).