

COGNITIVE
NEUROPSYCHOLOGY



**What lies beneath: A comparison of reading aloud in Pure
Alexia and Semantic Dementia**

Journal:	<i>Cognitive Neuropsychology</i>
Manuscript ID:	CN-PA 13-05.R2
Manuscript Type:	Full Article
Keywords:	Reading Aloud, Pure Alexia, Letter-by-Letter Reading, Semantic Dementia, Surface Dyslexia

SCHOLARONE™
Manuscripts

ABSTRACT

Exaggerated effects of word length upon reading aloud performance define Pure Alexia, but have also been observed in Semantic Dementia. Some researchers have proposed a reading-specific account, whereby performance in these two disorders reflects the same cause: impaired orthographic processing. In contrast, according to the primary systems view of acquired reading disorders, Pure Alexia results from a basic visual processing deficit, whereas degraded semantic knowledge undermines reading performance in Semantic Dementia. To explore the source of reading deficits in these two disorders, we compared the reading performance of 10 Pure Alexic and 10 Semantic Dementia patients, matched in terms of overall severity of reading deficit. The results revealed comparable frequency effects on reading accuracy, but weaker effects of regularity in Pure Alexia than Semantic Dementia. Analysis of error types revealed a higher rate of letter-based errors and a lower rate of regularisation responses in Pure Alexia than Semantic Dementia. Error responses were most often words in Pure Alexia but most often nonwords in Semantic Dementia. Although all patients made some letter substitution errors, these were characterised by visual similarity in Pure Alexia and phonological similarity in Semantic Dementia. Overall, the data indicate that the reading deficits in Pure Alexia and Semantic Dementia arise from impairments of visual processing and knowledge of word meaning, respectively. The locus and mechanisms of these impairments are placed within the context of current connectionist models of reading.

Keywords: Reading Aloud; Pure Alexia; Letter-by-Letter Reading; Semantic Dementia; Surface Dyslexia.

1
2
3 Efficient activation and integration of orthographic knowledge is essential in fluent
4 reading. Any disruption to this process as a consequence of brain damage will result in some
5 form of reading deficit, or acquired dyslexia. One such disorder is pure alexia (PA), which is
6 seen after damage to or disconnection of the left ventral occipito-temporal cortex (vOTC).
7 Behaviourally, the traditional definition of PA is as a highly selective reading deficit, without
8 associated problems in spoken language (aphasia), spelling (dysgraphia) or object recognition
9 (agnosia) (Déjerine, 1892). PA patients experience difficulties in accurate and rapid parallel
10 activation of the letters in words, which undermines their reading. This is evident in a very
11 marked effect of the number of letters in a word on patients' reading speed (Behrmann &
12 Plaut, 2013a; Roberts et al., 2013), which stands in contrast to the minimal effects of word
13 length seen in normal individuals' reading aloud (Henderson, 1982; Weekes, 1997). This
14 exaggerated length effect in PA is interpreted as reflecting sequential letter identification, or
15 letter-by-letter reading, and indeed some patients show this reading strategy overtly.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 While the hallmark length effect that defines PA is well established and accepted, the
33 cognitive cause of the reading deficit has been the matter of considerable debate. By one
34 account, PA is a reading specific disorder, and reports of patients who have shown normal
35 visual processing and recognition of objects have been used to support such a view (e.g., (Kay
36 & Hanley, 1991; Miozzo & Caramazza, 1998) and vice versa (e.g., (Yong, Warren,
37 Warrington, & Crutch, 2013). Within this approach, a number of researchers have suggested
38 that PA arises as a result of damage to an orthographic input lexicon (or its input connections)
39 (e.g., (Marshall & Newcombe, 1973; Noble, Glosser, & Grossman, 2000; Warrington &
40 Langdon, 1994; Warrington & Shallice, 1979; Warrington & Shallice, 1980), which contains
41 entries for all known word forms, and has been associated with left vOTC (Cohen et al.,
42 2002; Vinckier et al., 2007). As a result, these patients can no longer efficiently activate
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 word forms, so the letter-by-letter reading strategy functions to boost activation of
4
5 appropriate candidate lexical entries.
6
7

8 Reading-specific accounts that focus on damage to orthographic *lexical*
9
10 representations should predict an increased incidence of nonlexical reading responses which,
11
12 in the case of irregular words, would take the form of regularisation errors (e.g., sew read as
13
14 “sue”). While PA patients do show some evidence of enhanced effects of regularity on
15
16 reading aloud (Behrmann, Nelson, & Sekuler, 1998; Rapcsak & Beeson, 2004), regularisation
17
18 responses are relatively rare (Cumming, Patterson, Verfaellie, & Graham, 2006; Patterson &
19
20 Kay, 1982). Hence a different form of a reading-specific account proposed that PA patients
21
22 may in fact have difficulties with letter recognition, which would compromise input to both
23
24 lexical and nonlexical processing (Arguin & Bub, 1993; Behrmann & Shallice, 1995; Bub,
25
26 Black, & Howell, 1989; Hanley & Kay, 1996; Howard, 1991; Patterson & Kay, 1982; Perri,
27
28 Bartolomeo, & Silveri, 1996; Reuter-Lorenz & Brunn, 1990; Rosazza, Appollonio, Isella, &
29
30 Shallice, 2007). This account is consistent with the observation that PA patients often mis-
31
32 identify the component letters of words (Cumming, et al., 2006).
33
34
35
36
37

38 A contrasting perspective on PA is that it arises from a particular kind of visual deficit
39
40 that undermines the input to the reading system (Behrmann, Plaut, & Nelson, 1998; Farah &
41
42 Wallace, 1991). This view falls within the Primary Systems account of acquired dyslexia,
43
44 whereby reading disorders arise due to disruption of more basic visual, phonological and
45
46 semantic processing (Patterson & Lambon Ralph, 1999), which has been implemented in
47
48 connectionist models of reading (Chang, Furber, & Welbourne, 2012a; Plaut & Behrmann,
49
50 2011; Welbourne, Woollams, Crisp, & Lambon Ralph, 2011; Woollams, Lambon Ralph,
51
52 Plaut, & Patterson, 2007). Neuroimaging studies reveal that vOTC receives high acuity
53
54 foveal visual input (Hasson, Harel, Levy, & Malach, 2003; Hasson, Levy, Behrmann,
55
56 Hendler, & Malach, 2002; Levy, Hasson, Avidan, Hendler, & Malach, 2001; Malach, Levy,
57
58
59
60

1
2
3 & Hasson, 2002; Woodhead, Wise, Sereno, & Leech, 2011), which is particularly salient
4
5 when dealing with complex and confusable visual stimuli like letter strings. In line with this
6
7 view, pure alexia patients show reduced sensitivity to higher spatial frequency information
8
9 (Roberts, et al., 2013), although this is not universal (Starrfelt, Nielsen, Habekost, &
10
11 Andersen, 2013). Also in keeping with a visual deficit account, the exaggerated length effect
12
13 is accompanied by increased sensitivity to the visual confusability of letters (Arguin, Fiset, &
14
15 Bub, 2002; Fiset, Arguin, Bub, Humphreys, & Riddoch, 2005; Harris, Olson, & Humphreys,
16
17 2013; Johnson & Rayner, 2007). Interestingly, when higher spatial frequencies are
18
19 artificially removed, normal individuals show increased effects both of word length and letter
20
21 confusability (Fiset, Arguin, & Fiset, 2006; Tadros, Fiset, Gosselin, & Arguin, 2009). Yet
22
23 letter strings are by no means the only stimuli that rely on such information, with this same
24
25 brain region activated in face and object recognition (Behrmann & Plaut, 2013b; Malach, et
26
27 al., 2002; Nestor, Behrmann, & Plaut, 2013; Price & Devlin, 2003, 2011; Vogel, Petersen, &
28
29 Schlaggar, 2012; Woodhead, et al., 2011). By this account then, patients with damage to left
30
31 vOTC should show impairments in processing any visual stimuli that require medium to high
32
33 acuity foveal input for effective recognition.
34
35
36
37
38

39
40 When it has been assessed, the accuracy of non-linguistic visual processing in PA has
41
42 varied across cases, with some patients apparently showing normal performance (e.g.,(Kay &
43
44 Hanley, 1991; Miozzo & Caramazza, 1998), while others have shown significant
45
46 impairments (e.g.,(Cumming, et al., 2006; Roberts, et al., 2013). In studies that have also
47
48 considered reaction times, which is of course the measure by which their reading deficit is
49
50 defined, clear evidence of visual processing impairments has emerged, particularly for
51
52 complex stimuli. Behrmann et al. (1998a) reported five pure alexia patients to be slowed in
53
54 naming pictures, but only those high in visual complexity. Similarly, a large case series of 21
55
56 PA patients revealed significantly impaired performance in matching checkerboard stimuli
57
58
59
60

1
2
3 and unfamiliar logographic characters, most markedly for complex items in the presence of
4
5 visually similar distracters (Roberts et al., 2013, see also (Mycroft, Behrmann, & Kay, 2009).
6
7 Moreover, performance for this condition was strongly related to the severity of the reading
8
9 deficit, as measured by the size of the length effect.
10

11
12 Despite mounting evidence for a visual deficit in PA, this is unlikely to be the only
13
14 possible cause of abnormal word length effects, as these have also been reported in other
15
16 neuropsychological conditions, such as semantic dementia (SD) (Cumming, et al., 2006;
17
18 Gold et al., 2005; Patterson & Hodges, 1992). SD is a selective and progressive disorder of
19
20 conceptual knowledge associated with atrophy and hypometabolism of the anterior temporal
21
22 lobes (Adlam et al., 2006; Nestor, Fryer, & Hodges, 2006). Reading aloud in SD shows a
23
24 near-universal pattern of surface dyslexia, where words with exceptional spelling sound
25
26 correspondences, particularly those low in frequency, are read aloud according to more
27
28 typical correspondences (regularised). Moreover, accuracy for these exception items is
29
30 strongly related to the extent of the patients' receptive and expressive semantic deficits
31
32 (Graham, Patterson, & Hodges, 2000; Patterson et al., 2006; Woollams, et al., 2007). The
33
34 primary systems interpretation of these findings is that whole word semantic knowledge
35
36 supports the pronunciation of exception word items (Patterson & Lambon Ralph, 1999;
37
38 Patterson, et al., 2006).
39
40
41
42
43
44

45 Yet there have been a few reports of SD patients with accuracy of low-frequency
46
47 exception word reading falling within the normal range despite an appreciable semantic
48
49 deficit (Blazely, Coltheart, & Casey, 2005; Cipolotti & Warrington, 1995). This has led
50
51 some researchers to propose that exception word reading in SD is undermined not by
52
53 semantic deficits associated with ATL damage, but rather the posterior spread of atrophy into
54
55 the left vOTC region (Coltheart, Tree, & Saunders, 2010). This account predicts that there
56
57 should be clear similarities in the reading aloud performance of SD and PA patients. The
58
59
60

1
2
3 observation of abnormally strong length effects in SD (Cumming, et al., 2006; Gold, et al.,
4
5 2005), combined with reports of SD cases who have adopted an explicit letter-by-letter
6
7 reading strategy (Noble, et al., 2000), have been considered evidence for this view. An
8
9 alternative perspective, however, is that it is these length effects arise as a consequence of
10
11 reduced support from whole word semantic knowledge that would usually bind the letters of
12
13 a word together, offsetting costs associated with more letters.
14
15

16
17 In a direct comparison of the visual processing and reading performance of 3 PA
18
19 patients with 3 SD patients (Cumming, et al., 2006), performance on non-verbal visual
20
21 processing tasks for both familiar and unfamiliar objects was normal in SD, but impaired in
22
23 PA. Letter matching was normal for SD at longer durations, whereas in PA it was universally
24
25 impaired. Length effects were seen in both types of disorder, but these were significantly
26
27 smaller for the SD than PA patients (although it should be kept in mind that accuracy was
28
29 higher in SD than PA). Interestingly, error responses were usually words for the PA patients,
30
31 but nonwords for the SD patients. This is consistent with work showing enhanced influences
32
33 of whole word variables in PA (e.g.,(Roberts, Lambon Ralph, & Woollams, 2010). The
34
35 notion of a bottom-up visual and a top-down semantic impairment both increasing length
36
37 effects was reinforced by the finding that PA patients showed smaller length effects for words
38
39 than nonwords, while SD patients showed equivalent effects. Taken together, these results
40
41 speak to a visual origin of length effects in PA and a semantic cause in SD.
42
43
44
45

46
47 The goal of the present research was to illuminate the source of reading deficits in PA
48
49 and SD by comparing patients matched on overall severity. Previous work has already
50
51 compared the effects of length and lexicality in PA and SD (Cumming, et al., 2006), so here
52
53 we explored the impacts of frequency and regularity using the Surface List (Patterson &
54
55 Hodges, 1992) and considered not only overall accuracy but also the nature of the patients'
56
57 reading errors. If the deficits in both PA and SD arise from damage to reading-specific
58
59
60

1
2
3 orthographic processing, we would expect to see similar reading performance across the two
4
5 groups. If, in contrast, the two reading deficits arise from underlying visual and semantic
6
7 causes, respectively, then we would expect (a) weaker effects of regularity for PA than SD,
8
9 (b) a higher proportion of nonword and regularisation responses in SD than PA, and (c) a
10
11 higher proportion of incorrect word responses and letter-based errors in PA than SD.
12
13

14 15 16 17 18 **Method:**

19 20 21 *Participants:*

22
23
24 Pure Alexia: For this study we operationally characterised Pure Alexia in terms of a
25
26 combination of damage to the left occipito-temporal cortex combined with slowed reading
27
28 and an abnormally large word length effect. Ten PA patients with overt LBL reading of
29
30 varying degrees participated. All were native speakers of English who had suffered from
31
32 acute brain injury more than two years prior to the time of testing. These patients were
33
34 recruited from local NHS speech and language therapy services on the basis of marked
35
36 increases in word-reading latency as a function of letter length. On our reading list of 180
37
38 words (Roberts, et al., 2010), overt LBL responses were produced by every patient.
39
40
41

42
43 As can be seen in Table 1, all patients had damage in the occipito-temporal region, as
44
45 judged by a neurologist, as a consequence of stroke or tumour resection. Scans for eight of
46
47 the ten patients are provided in the Appendix. Scans for two other patients (PM and KW)
48
49 were not available, hence the determination of damage was made on the basis of the
50
51 neurologist's written report. Overall, neuropsychological background assessment indicated
52
53 that the patients had preserved working memory (digit span (Wechsler, 1987)) and
54
55 phonological processing, with only a one patient slightly impaired on the more demanding
56
57
58
59
60

1
2
3 tests of phonological segmentation (EW). Deficits in visual processing on at least one subtest
4
5 of the VOSP (Warrington & James, 1991) were apparent in all patients.
6
7

8 Performance on the Cambridge Picture Naming test (Bozeat, Lambon Ralph,
9
10 Patterson, Garrard, & Hodges, 2000) revealed impaired performance in all bar one case (PM).
11
12 Receptive semantic processing tests included the Cambridge Spoken Word to Picture
13
14 Matching test (Bozeat, Lambon Ralph, et al., 2000) where a spoken word was matched to a
15
16 target picture amongst nine semantically related alternatives; the Camel and Cactus Pictures
17
18 test (Bozeat, Lambon Ralph, et al., 2000) , where a target picture was matched to a picture of
19
20 an associated item in the context of three semantically similar items; and the 96 Synonyms
21
22 test (Jefferies, Patterson, Jones, & Lambon Ralph, 2009), where a written target word was
23
24 matched to a synonym in the context of two other related words of similar frequency and
25
26 imageability (options are also read to the patient by the experimenter). Six patients (TS, KW,
27
28 SC, EW, MS, & AT) showed mild but measureable impairments on at least one of these
29
30 receptive semantic tests.
31
32
33
34
35

36 While the prevalence of deficits on these semantic tests could be interpreted as
37
38 indicating deficits in conceptual knowledge, it is worth noting that all of the tests involved
39
40 either pictures or written words. Poor performance on these tests is therefore consistent with
41
42 optic aphasia, if conceptualised as a disconnection of semantics from visual input (Plaut &
43
44 Shallice, 1993). Yet in light of the demonstrated visual impairments on the VOSP, it seems
45
46 plausible that impaired performance on the semantic tests in this patient group may have
47
48 arisen as a consequence of problems in visual processing. We hypothesise that reduced
49
50 sensitivity to higher spatial frequencies could impair performance on (a) the more demanding
51
52 subtests of the VOSP such as progressive silhouettes; (b) semantic tests that involve picture
53
54 identification; and (c) semantic tests that also involve reading written words. Such an
55
56 account would of course be consistent with the primary systems view and previous reports of
57
58
59
60

1
2
3 object processing deficits in this population (e.g. (Behrmann, Nelson, et al., 1998; Mycroft, et
4
5 al., 2009; Roberts, et al., 2013).

6
7
8 Data for spelling words of different lengths from the PALPA 39 subtest (Kay, Lesser,
9
10 & Coltheart, 1992b) were available for only five cases. Although this test does not have
11
12 published norms, according to <http://www.neuro.mcw.edu/mcword/>, the mean frequency of
13
14 items is 108 per million, and the control range on the PALPA 40 is between 30 and 100% for
15
16 items with a mean frequency 105 per million. Hence, although spelling performance was not
17
18 perfect in all cases, it would seem there was good performance for shorter words of three or
19
20 four letters, and performance for longer words was also good in most cases, with the only
21
22 clearly impaired case being EW. **Given that EW also showed deficits in tests of phonological
23
24 and semantic processing, it is possible that aphasic deficits may have contributed to her
25
26 impaired orthographic processing.**

27
28
29 All patients showed elevated mean reading speeds on the 180-item list from Roberts
30
31 et al. (2010) (RTs were derived using a voice recorder and manual analysis of reaction time
32
33 data using WavePad software). All patients showed an appreciable influence of word length
34
35 upon their reading speed, although the strength of the effect varied across different patients.
36
37 This variability is also reflected in accuracy of Surface List reading, and demonstrates that
38
39 any comparisons across patient types must take into account overall severity of the reading
40
41 disorder.

42
43
44 Semantic Dementia: Ten SD patients with reading accuracy comparable to that of the
45
46 PA patients on high-frequency regular words were selected from the cohort presented in
47
48 Woollams et al. (2007). All patients had received a diagnosis of Semantic Dementia
49
50 according to the (Neary et al., 1998) consensus criteria, which include atrophy of the anterior
51
52 temporal lobe. Their selective semantic impairment is apparent in Table 2.
53
54
55
56
57
58
59
60

1
2
3 MMSE Scores (Folstein, Folstein, & McHugh, 1975) were below the control range
4
5 for all patients, as would be expected given that this test assesses some aspects of verbal
6
7 ability. Working memory performance as assessed by digit span (Wechsler, 1987) was
8
9 within the normal range in all but one case (DA1). Visuo-perceptual processing was
10
11 reasonably intact, as indicated by scores within the normal range for all patients on the Rey
12
13 Immediate Copy Test (Lezak, 1976). Where available, data from the VOSP showed
14
15 preserved performance except for the Silhouettes subtest and in one case (MB1) on the
16
17 Object Decision subtest, which is understandable given this draws on knowledge of object
18
19 identity.
20
21
22
23

24 There was a marked impairment on tests tapping semantic memory. Performance was
25
26 outside the control range for all patients on both the Cambridge Picture Naming and Spoken
27
28 WPM tests (Bozeat, Lambon Ralph, et al., 2000; Hodges, Patterson, Oxbury, & Funnell,
29
30 1992), and on the Pyramids and Palm Trees Test (Howard & Patterson, 1992), reflecting the
31
32 progressive anomia and declining comprehension that are key features of SD. Deficits in
33
34 semantically-generated output are apparent on the Category Fluency Test (Hodges, Salmon,
35
36 & Butters, 1992), in which patients are asked to generate as many examples as they can in
37
38 one minute each for eight semantic categories, arguing against a visual contribution to the
39
40 decreased performance seen on the semantic tests. Performance on the Surface List shows a
41
42 consistent pattern of Surface Dyslexia, with all patients showing poor performance for low
43
44 frequency exception words.
45
46
47
48

49 *Stimuli:*

50
51
52 The reading performance of all PA and SD patients was assessed using the Surface
53
54 List (Patterson & Hodges, 1992, see Woollams et al., 2007 Appendix A). The Surface List
55
56 consists of a factorial manipulation of frequency and regularity, with 42 items per cell.
57
58
59
60

1
2
3 Within each level of frequency, the regular and exception items are matched on initial
4 phoneme, and do not differ according to (Kučera & Francis, 1967) written frequency
5 (HFR=811.43, HFE=798.83, $t(1,80) < 1$; LFR=5.78, LFR=5.41, $t(1,78) < 1$) or orthographic
6 length (HFR=4.14, HFE=4.24, $t(1,82) < 1$; LFR=4.83, LFR=4.81, $t(1,82) < 1$).
7
8
9
10

11
12
13 *Procedure:*

14
15 For the PA patients, after an initial series of 12 practice items, patients viewed each
16 item of the Surface List one at a time in the centre of a laptop screen. Items were displayed
17 using e-prime software (Schneider, Eschman, & Zuccolotto, 2002) with an input of Arial 18
18 point that translated to the equivalent of 34 point once displayed on the screen
19 (ascender/descender height = 0.9cm). Responses were digitally recorded for later coding.
20 For the SD patients, practice and test items were presented one at a time on cards in Geneva
21 26 point font (ascender/descender height = 0.7cm), and responses were coded in written form
22 by the experimenter. Note that although presentation format differed over patient group, the
23 two are near identical proportional fonts (e.g., pint vs pint), and while the font size was larger
24 for the PA patients than the SD patients, this in fact works against our hypothesis of a more
25 visual errors for PA than SD patients, as letter identification has been shown to be relatively
26 independent of such variations in size (Pelli, Burns, Farell, & Moore-Page, 2006). For both
27 groups, test items were presented in a fixed pseudo-random order that ensured a
28 representative distribution of items from each condition over four blocks.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **Results:**

49
50 *Accuracy:*

51
52 Reading accuracy for the PA and SD patients is presented in Figure 1. Data were
53 analysed using a 2 (Patient: PA/SD) by 2 (Frequency: High/Low) by 2 (Regularity:
54
55
56
57
58
59
60

Regular/Exception) ANOVA, with repeated measures on the latter two factors. The results revealed no main effect of patient type ($F(1,18)=0.74$, $p=.401$), indicating that the severity matching had been successful. There were significant main effects of both frequency ($F(1,18)=41.25$, $p<.0005$) and regularity ($F(1,18)=28.55$, $p<.0005$), and their interaction ($F(1,18)=12.49$, $p=.002$). The two patient types showed comparable effects of frequency ($F(1,18)=1.85$, $p=.191$), but the impact of regularity was significantly stronger in SD than PA ($F(1,18)=8.95$, $p=.008$). The significant three way interaction ($F(1,18)=6.27$, $p=.022$) was driven by the SD patients' significantly worse performance specifically on the low frequency exception words ($t(18)=2.49$, $p=.011$, one-tailed). Repeated measures ANOVAs on the PA patients alone showed significant main effects of frequency ($F(1,9)=19.10$, $p=.002$), a marginal effect of regularity ($F(1,9)=4.51$, $p=.063$), and no interaction between them ($F(1,9)=0.38$, $p=.551$). A parallel analysis on the SD patients alone showed significant main effects of frequency ($F(1,9)=22.79$, $p=.001$), regularity ($F(1,9)=25.05$, $p=.001$), and an interaction between them ($F(1,9)=29.72$, $p<.0005$).

Error types:

All errors were transcribed in order to maximise orthographic similarity to the target. A variety of error types were observed amongst both PA and SD patients, and a summary of these is provided in Table 3. We classified each error into one of the following mutually exclusive categories (i) omissions (which were rare in both groups); (ii) Legitimate Alternative Reading of Components (LARC), in which the patient pronounced the word in line with spelling-sound correspondences of one or more other known words (e.g., *sew* > "*sue*", as in *few* and *stew*); (iii) visual errors, in which the response had at least 1 letter (out of 3 or 4) or 2 letters (out of 5 or 6) in common with target (e.g., *saw* > "*save*"; *cough* > "*coach*"); (iv) letter omissions, where all letters of the response were found in the target, but the response was one letter shorter than the target (e.g., *learn* > "*lean*"); (v) letter

1
2
3 additions, where all letters of the response were found in the target, but the response was one
4
5 letter longer than the target (e.g., *per* > “*pear*”); (vi) letter transpositions, where the
6
7 response was the same length and contained all the letters of the target, but two adjacent
8
9 letters had switched order (e.g., *trial* > “*trail*”); and (vii) letter substitutions, where the
10
11 response was the same length as the target but one letter had been replaced (e.g., *food* >
12
13 “*fool*”).
14
15

16
17 As can be seen in Table 3, omission errors were very rare in the PA patients, but as
18
19 they were non-existent in the SD patients, this group difference was significant ($t(18)=1.97$,
20
21 $p=.032$, one-tailed). As expected LARC errors were the most prevalent type of error for the
22
23 SD patients and, while some LARC errors were made by the PA patients, these were
24
25 significantly less common ($t(18)=3.21$, $p=.002$, one-tailed). Visual errors were marginally
26
27 more common for the PA than SD patients ($t(18)=1.47$, $p=.079$, one-tailed). Neither letter
28
29 omissions nor additions differed significantly between PA and SD patients ($t(18)=1.25$,
30
31 $p=.115$, one-tailed; $t(18)=0.89$, $p=.194$, one-tailed, respectively). Letter transpositions,
32
33 although rare overall, were significantly more common in PA than SD patients ($t(18)=2.16$,
34
35 $p=.022$, one-tailed). The most prevalent error type for the PA patients was letter
36
37 substitutions, and while such errors were also seen in the SD patients, they were significantly
38
39 less common ($t(18)=2.42$, $p=.013$, one-tailed). No difference between PA and SD patients on
40
41 other error types was apparent ($t(18)=0.35$, $p=.364$, one-tailed). To summarise, LARC errors
42
43 were significantly more common for the SD than the PA patients, whereas at least some types
44
45 of letter-based errors (visually related responses, transpositions and substitutions) were
46
47 significantly more common in the PA than the SD patients. This pattern is displayed in
48
49 Figure 2, and is consistent with reading performance disrupted by a semantic deficit in SD
50
51 and by a visual deficit in PA.
52
53
54
55
56
57
58
59
60

1
2
3 For all errors of commission, we also coded whether the responses corresponded to
4 another known word, and these proportions are displayed in Table 4. There was a highly
5 significant difference between the PA and SD patients on this measure ($t(18)=6.87$,
6 $p<.000005$, one-tailed). As can be seen in Figure 3, the vast majority of errors of commission
7 produced by the PA patients were words. The SD patients, on the other hand, were
8 somewhat more likely to produce nonword than word errors. This striking difference is
9 consistent with the idea that reading aloud in SD is characterised by a reduction of semantic
10 activation, such that there is insufficient top-down information to prevent nonword responses.
11 In contrast, the high proportion of word errors in the PA patients suggests that reading
12 responses in the face of compromised bottom-up visual input are typically constrained by
13 top-down information.

24
25
26
27
28 *Letter substitutions:*

29
30
31 The analysis of error types demonstrated that PA patients were significantly more
32 likely than SD patients to substitute one of the component letters of a word. If these
33 substitutions result from a visual processing deficit in PA, then we would also expect that the
34 substitutions result from a visual processing deficit in PA, then we would also expect that the
35 form of these errors will be driven more by visual similarity than in SD. To assess this
36 hypothesis, we coded the letter presented and letter 'reported' (as reflected in the whole
37 response) according to the letter confusability matrix in Patterson and Kay (1982), derived
38 from the errors made by normal participants in identifying letters presented briefly in
39 peripheral vision (Bouma, 1971). We selected this confusability matrix because: (i) it was
40 based on lower-case letters, as used in our reading list; (ii) it was derived from peripheral
41 vision, resulting in perception with reduced medium to high spatial frequency information,
42 akin to deficits suggested in PA patients (Roberts, et al., 2013); and (iii) it has been used
43 before with reference to letter substitutions in cases of PA (Patterson & Kay, 1982). The
44 results for each group can be seen in Figure 4, where the values represent the proportion of all
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 substitutions. The cells closest to the diagonal represent maximum visual similarity, and the
4
5 substitutions of the PA patients fall closer to the diagonal than the SD patients, as
6
7 hypothesised. In order to quantify this difference, we computed the Euclidean distance
8
9 between the presented and reported letters within the matrix for each error in the following
10
11 way: we created a matrix where each letter was assigned a number from 1 to 26 (e.g., a=2,
12
13 o=3), and the absolute difference between the presented and reported letter yielded a distance
14
15 for that confusion for a given patient (e.g., cat read as “cot” had a distance of 1). The average
16
17 distance for PA patients across all letter substitution errors was 5.1, while that for SD patients
18
19 was 7.0, which was significantly lower ($t(165)=2.38$, $p=.009$, one-tailed). This result is again
20
21 consistent with a visual deficit undermining reading in PA.
22
23
24
25

26 The preceding analysis indicates a key role for visual similarity in the specific letter
27
28 substitution errors of the PA patients. What might be the relevant relationship between
29
30 stimulus and response words in SD reading errors? One possibility is that semantic
31
32 impairment exerts its effects on reading aloud through mild perturbation of
33
34 phonological/phonetic processing. To assess this hypothesis, we used (Bailey & Hahn, 2005)
35
36 coding scheme to capture the sound similarity - in terms of number of shared features (place,
37
38 manner, voice, sonorance) - corresponding to the phonemes involved in letter substitution
39
40 errors. The results can be seen in Figure 5, where the values represent the proportion of all
41
42 consonant-consonant substitutions. This reveals that the SD patients' letter substitutions were
43
44 more likely to equate to phonemes sharing 2 or 3 features with the target phoneme, whereas
45
46 for PA patients such substitutions typically shared either no, or just a single, phonetic feature.
47
48 A comparison of the average number of shared phonetic features demonstrated greater
49
50 phonemic similarity of substitutions amongst the SD (2.53) than PA patients (2.14)
51
52 ($t(165)=1.69$, $p=.038$, one-tailed).
53
54
55
56
57

58 Discussion:

1
2
3 This study investigated the extent to which the reading deficits seen in PA and SD
4 arise from similar or different causes. The fact that increased word length effects have been
5 seen in reading performance in both these disorders has led some researchers to propose that
6 they share a common cause in terms of disruption to reading-specific orthographic processing
7 (e.g.,(Coltheart, et al., 2010; Noble, et al., 2000). In contrast, the primary systems view
8 attributes all characteristics of these two reading disorders, including length effects, to a
9 deficit in general visual processing in PA and to a deficit in central semantic processing in SD
10 (Patterson & Lambon Ralph, 1999; Roberts, et al., 2013; Woollams, et al., 2007). These two
11 accounts therefore diverge in the extent to which they predict resemblances between reading
12 performance in the two disorders. Here, we explored this issue by directly comparing the
13 impact of frequency and regularity on reading accuracy, and the nature of error types, in 10
14 cases of PA and 10 cases of SD who were matched on their accuracy in reading single words
15 aloud.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 In terms of reading accuracy, the PA and SD patients were similar in that they showed
33 comparable effects of frequency, which concurs with results previously reported in the
34 literature (e.g.,(Behrmann, Plaut, et al., 1998; Graham, et al., 2000). While this result could
35 be consistent with a shared locus of impairment in orthographic processing, it could also arise
36 from different sources. In PA, the perception of high frequency words may be less disrupted
37 due to feedback from intact higher-order linguistic/semantic representations (Roberts, et al.,
38 2010), whereas in SD, the production of low frequency words may be more disrupted because
39 semantic representations of these items are most vulnerable to damage (Lambon Ralph,
40 Graham, Ellis, & Hodges, 1998; Rogers et al., 2004; Woollams, Cooper-Pye, Hodges, &
41 Patterson, 2008). This notion that the influence of top-down activation is increased in PA but
42 reduced in SD is consistent with the striking finding reported here and previously (Cumming,
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 et al., 2006) that PA patients are much more likely to produce errors that are nevertheless
4
5 known words, while SD patients are in fact more likely to produce errors that are nonwords.
6
7

8 The impact of regularity on reading accuracy was significantly weaker in PA than SD,
9
10 and the incidence of LARC errors was also significantly lower. Consistent with previous
11
12 work, there was a marginally significant effect of regularity on PA reading accuracy
13
14 (Behrmann, Nelson, et al., 1998; Rapcsak & Beeson, 2004), but LARC errors were the least
15
16 common error type for the PA patients (Cumming, et al., 2006; Patterson & Kay, 1982). This
17
18 contrasts with the very strong impact of regularity on reading accuracy for the SD patients,
19
20 and the fact that LARC errors were the most common error type in SD, as has been
21
22 previously seen in larger samples (Graham, et al., 2000; Woollams, et al., 2007). The
23
24 prevalence of LARC errors in SD speaks to intact processing along a direct pathway between
25
26 orthography and phonology in the face of compromised whole-word knowledge due to
27
28 damage to the semantic system.
29
30
31
32

33 Consideration of the nature of reading errors also highlighted a higher incidence of
34
35 certain letter based errors in PA than SD – specifically those where the stimulus and response
36
37 shared most of their letters (visual errors, see also Rapcsak & Beeson, 2004), where letters in
38
39 the response re-ordered those in the stimulus (transpositions: see also (Pflugshaupt et al.,
40
41 2011), and where a single letter in the stimulus was replaced by another in the response
42
43 (substitutions, see Patterson & Kay, 1982). Indeed, it was letter substitutions that were the
44
45 most common type of error for the PA patients, but some substitution errors were also
46
47 produced by the SD patients. To understand the source of the substitution errors in the two
48
49 patient types, we first considered the extent to which the presented and reported letters were
50
51 visually similar, as measured by their degree of confusability by normal participants when
52
53 letters are presented in peripheral vision (Bouma, 1971; Patterson & Kay, 1982), a technique
54
55 which may simulate the lower spatial frequency information available to PA patients with
56
57
58
59
60

1
2
3 unlimited duration central presentation (Roberts, et al., 2013). The visual similarity of the
4
5 presented and reported letters was significantly higher in PA than SD, consistent with a visual
6
7 processing impairment as the cause of the reading deficit in PA.
8
9

10 We then further explored the source of letter substitution errors in SD by considering
11
12 the extent to which they were driven by phonological similarity, as measured by overlap in
13
14 terms of the phonetic features of the presented and reported consonant phonemes (Bailey &
15
16 Hahn, 2005). The motivation behind this analysis was the possibility that semantic damage
17
18 could exert its effects on reading aloud through disruption of phonological processing. This
19
20 notion is supported by a body of literature demonstrating poorer repetition by SD patients of
21
22 short sequences of words whose meanings they no longer know, when compared to words
23
24 with meanings that are still known (Knott, Patterson, & Hodges, 1997, 2000; Patterson,
25
26 Graham, & Hodges, 1994). This poorer performance is characterised by phoneme migration
27
28 errors (e.g., mint, rug will be reproduced as rint, mug), suggesting that semantic activation
29
30 helps to bind together phonological elements. Consistent with this view, the phonological
31
32 similarity of the letter substitutions of SD patients was significantly higher than for PA
33
34 patients.
35
36
37
38
39

40 The phonological similarity of letter substitution errors in SD does suggest that
41
42 semantic impairment exerts effects on reading aloud through disruption of phonological
43
44 processing, but there are multiple mechanisms by which this could occur. SD patients' poor
45
46 performance in repetition of lists of words with degraded meaning has been viewed as
47
48 reflecting dramatically reduced semantic activation of phonology, consistent with the
49
50 prevalence of omission errors in SD patients' picture naming (Woollams, et al., 2008) and the
51
52 ineffectiveness of phonological cueing for their anomia (Graham, Patterson, & Hodges, 1995;
53
54 Jefferies, Patterson, & Lambon Ralph, 2008). It is possible that degraded knowledge not only
55
56 reduces phonological activation but also adds noise to it, consistent with the occurrence of
57
58
59
60

1
2
3 errors of commission in SD picture naming (Woollams, et al., 2008). This noisy activation
4
5 would be inherited by phonological representations during reading, and indeed this is the
6
7 approach taken by Woollams et al. (2007) in their simulations of reading aloud in SD within
8
9 the connectionist triangle model of (Plaut, McClelland, Seidenberg, & Patterson, 1996). To
10
11 the extent that phonological representations are organised according to phonetic features
12
13 (e.g.,(Harm & Seidenberg, 2004), then this noisy activation would result in the substitution of
14
15 similar phonemes during reading aloud, as observed in the present study.
16
17
18

19
20 Overall then, a consideration of the reading aloud performance in PA and SD patients
21
22 matched for accuracy of reading aloud has shown that the two groups perform very
23
24 differently. The prevalence of visual errors and the visual similarity of letter substitutions in
25
26 PA indicate a general visual processing deficit, whereas the prevalence of LARC errors and
27
28 the phonetic similarity of phoneme substitutions in SD are consistent with a semantic
29
30 impairment, in line with a primary systems account of reading disorders. This account of PA
31
32 and SD reading is represented schematically in Figure 6 within the connectionist triangle
33
34 framework. The assumption of a general visual processing deficit in PA is supported not
35
36 only by the present data, but also previous work showing visual processing deficits to varying
37
38 degrees in these patients (Behrmann, Nelson, et al., 1998; Behrmann & Plaut, 2013a;
39
40 Behrmann & Shallice, 1995; Farah & McClelland, 1991; Friedman & Alexander, 1984;
41
42 Mycroft, et al., 2009; Roberts, et al., 2013; Starrfelt & Behrmann, 2011; Starrfelt, Habekost,
43
44 & Gerlach, 2010; Starrfelt, Habekost, & Leff, 2009), and recent neuroimaging work
45
46 implicating the vOTC in the processing of high spatial frequency foveal visual information
47
48 (Hasson, et al., 2003; Hasson, et al., 2002; Levy, et al., 2001; Malach, et al., 2002; Vogel, et
49
50 al., 2012; Woodhead, et al., 2011). The assumption of disruption specifically to semantics is
51
52 similarly supported by patient neuroimaging data: SD patients have structural and functional
53
54 abnormality of the ATL but not vOTC (Acosta-Cabronero et al., 2011; Nestor, et al., 2006;
55
56
57
58
59
60

1
2
3 Woollams, Lambon Ralph, Plaut, & Patterson, 2010), and the extent of ATL damage has
4
5 been directly linked to level of success on non-reading semantic tasks (Adlam, et al., 2006;
6
7 Mion et al., 2010).
8
9

10
11 Within the primary systems account, the visual deficit in PA undermines input to
12
13 orthographic processing, producing the patients' visual errors, letter transpositions and
14
15 visually similar letter substitutions. This can, however, be offset to some extent by top-down
16
17 activation from intact semantic and phonological information feeding back to orthography,
18
19 producing the effects of frequency (and possibly regularity) observed here, combined with
20
21 the prevalence of real-word error responses. In contrast, the semantic impairment in SD
22
23 reduces and disrupts activation of phonology during reading, increasing the incidence of
24
25 nonword error responses. Effects of frequency arise because semantic representations of low
26
27 frequency words are less robust to damage, while regularity effects arise as reading of words
28
29 with atypical spelling-sound mappings come to rely more upon semantic activation of
30
31 phonology over the course of learning (Plaut, et al., 1996). The intact mappings directly
32
33 between orthography and phonology produce LARC errors in the case of words with atypical
34
35 mappings, particularly those low in frequency. In some cases the direct activation of
36
37 phonology can be disrupted by the noise from degraded semantic activations, and the result is
38
39 the substitution of a phonetically similar phoneme.
40
41
42
43

44
45 Our account requires further exploration within implemented connectionist
46
47 computational models of reading aloud. Some of these models incorporate phonological
48
49 representations in the form of phonetic features (e.g.,(Harm & Seidenberg, 2004), allowing
50
51 exploration of SD patients' errors. More recently, connectionist models have been extended
52
53 to accept raw visual input (Chang, et al., 2012a; Chang, Furber, & Welbourne, 2012b) and
54
55 could therefore potentially simulate PA patients' reading behaviour. This investigation has
56
57 provided target data for such simulations, and has demonstrated that despite surface
58
59
60


1
2
3 similarities in the reading impairments of PA and SD patients, a deeper consideration
4
5 indicates that these arise due to distinct impairments of visual processing vs. semantic
6
7 representation.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table 1. *Demographic and background neuropsychological data for the ten pure alexic patients included in the current study, ordered from least to most impaired according to high frequency regular word reading accuracy.*

	Max.	PM	JW	JM	TS	KW	SC	JWF	MS	AT	EW
<u>Demographics</u>											
Age	-	64	59	67	57	44	81	54	70	73	74
Years of education	-	10	11	10	10	10	11	10	10	10	10
<u>Lesion information</u>											
Neuroimaging summary	-	Occipito-temporal	Occipito-temporal	Occipito-temporal	Occipito-temporal	Occipito-temporal	Occipito-temporal	Occipito-temporal	Occipito-temporal	Occipito-parietal	Occipito-temporo-parietal
Aetiology	-	PCA stroke	PCA stroke	PCA tumour resection	PCA tumour resection	MCA stroke	PCA stroke	Post aneurism PCA infarct	PCA stroke	PCA stroke	MCA stroke
<u>Working memory</u>											
Digit span (scaled score)	18	NT	9	15	8	14	7	10	14	10	7
<u>Visual processing</u>											
Right visual field impairment*	-	Upper quadrant	Hemianopia	Upper quadrant	Hemianopia	Hemianopia	Hemianopia	Hemianopia	Hemianopia	Upper quadrant	Hemianopia
VOSP											
Incomplete letters	20	NT	19	20	19	20	18	17	16	16	19

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



Silhouettes	30	NT	25	18	22	19	3	24	19	13	12
Object decision	20	NT	17	17	18	20	14	19	16	17	17
Progressive silhouettes	20	NT	8	11	5	16	NT	NT	9	9	12
Dot counting	10	NT	10	10	10	9	10	10	9	10	10
Position discrimination	20	NT	20	20	18	20	17	16	19	20	20
Number location	10	NT	10	10	10	10	10	8	10	9	10
Cube analysis	10	NT	9	10	10	4	9	10	7	10	7
<u>Semantic processing</u>											
Naming ⁺	64	64	59	60	40	56	59	54	45	54	45
Spoken Word to Picture matching ⁺	64	NT	64	63	63	63	62	NT	62	63	57
Camel and Cactus (pictures) ⁺	64	64	52	61	24	NT	NT	61	47	NT	45
96 Synonyms (%) ⁻	96	NT	93	93	83	74	71	94	81	NT	76
<u>Phonological processing</u>											
PALPA 2: Phonological judgement	72	NT	71	72	68	71	NT	72	71	NT	65
PALPA 15: Rhyme judgement	60	NT	57	56	56	59	NT	58	53	NT	56
Phonological segmentation [#]	96	NT	96	94	87	97	NT	96	91	NT	69
<u>Spelling</u>											
PALPA 39 Written											
Short	100	NT	92	100	100	NT	NT	NT	100	NT	75

Long	100	NT	50	75	100	NT	NT	NT	75	NT	17
Reading Aloud											
180-item list:	-										
Mean RT (ms)	-	1013	7530	5432	5158	5903	7910	6484	12667	15683	7010
Mean Accuracy (%)	-	100	91	96	95	94	83	53	75	57	58
Length effect (ms per letter)	-	170	1299	911	1060	651	1843	1369	1650	523	2248
Length effect (% per letter) [^]	-	0	0.75	-1.25	-1.25	0	-3.75	1.5	-8.75	-4	-3.25
Surface List:											
High Frequency Regular	42	41	41	41	40	39	36	34	32	30	29
Low Frequency Regular	42	40	38	37	39	36	26	39	24	22	15
High Frequency Exception	42	39	40	39	37	41	31	37	30	29	25
Low Frequency Exception	42	37	28	33	34	32	16	34	20	27	26

Bold denotes abnormal performance represented by scores falling beyond two standard deviations below control performance where normative data available; ;For Digit Span, abnormal scores are two standard deviations below age appropriate means (Ivnik et al., 1992); PCA = Posterior Cerebral Artery; NT - Not tested; NA = Not available. * = Assessed using LERNREHA from Kasten, Strasburger, & Sabel (1997); VOSP: Visual Object and Space Perception battery (Warrington & James, 1991); PALPA: Psycholinguistic Assessment of Language Processing in Aphasia (Kay, Lesser, & Coltheart, 1992a); ⁺ Tests from (Bozeat, Gregory, Lambon Ralph, & Hodges, 2000); [~] Test from Jefferies et al. (2009); [#] Tests from (Patterson & Marcel, 1992); [^] represents decrease in accuracy for each additional letter in string..

Table 2. *Demographic and background neuropsychological data for the ten semantic dementia patients included in the current study, ordered from least to most impaired according to high frequency regular word reading accuracy.*

	Max.	GC6	LS3	MB1	DC1	DA1	AM4	NS2	MA6	FM8	AT6
<u>Demographics</u>											
Age	-	60	62	65	77	75	65	69	73	57	68
Years of education	-	12	13	11	8	16	16	9	13	10	19
<u>Cognitive Status</u>											
MMSE	-	NT	24	22	18	9	8	25	5	22	15
Raven's coloured	-	NT	NT	17	33	NT	30	36	NT	25	34
<u>Working memory</u>											
Digit span (scaled score)	18	5	7	4	7	3	6	6	NT	4	9
<u>Visual processing</u>											
Rey Immediate Copy	36	34	29	26	32	34	35	36	NT	32	36
VOSP											
Incomplete letters	20	20	19	19	18	17	NT	19	NT	NT	18
Silhouettes	30	5	NT	9	1	NT	NT	NT	NT	NT	NT
Object decision	20	14	NT	13	17	18	NT	NT	NT	NT	NT
Progressive silhouettes	20	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT
Dot counting	10	10	10	9	10	9	NT	10	NT	NT	10
Position discrimination	20	20	20	20	20	19	NT	20	NT	NT	20

5	Number location	10	8	10	7	10	10	NT	9	NT	NT	8
6	Cube analysis	10	10	9	8	10	8	NT	10	NT	NT	10
9	<u>Semantic processing</u>											
10	Naming ⁺	64	13	34	44	20	12	0	8	10	0	5
11	Spoken Word to Picture matching ⁺	64	35	60	58	44	50	17	42	43	59	29
12	Pyramids and Palm Trees (pictures)	52	NT	30	39	40	39	35	39	26	46	38
13	Pyramids and Palm Trees (words)	52	NT	NT	NT	30	41	30	33	26	NT	27
14	Category (8 categories)	-	NT	NT	25	13	4	1	19	0	0	NT
15	<u>Reading Aloud</u>											
16	Surface List:											
17	High Frequency Regular	42	41	41	40	39	39	36	36	33	32	29
18	Low Frequency Regular	42	40	35	41	39	37	34	15	25	29	29
19	High Frequency Exception	42	38	40	40	33	35	33	31	37	21	21
20	Low Frequency Exception	42	20	26	32	20	28	21	7	26	15	10

Bold denotes abnormal performance represented by scores falling beyond two standard deviations below performance of a group of between 100 and 24 (depending on the test) control participants comparable in terms of age and education; For Digit Span, abnormal scores are two standard deviations below age appropriate means (Ivnik, et al., 1992); For the Raven's, abnormal performance is that below the 50%th centile for older controls in norms ; NT - Not tested; VOSP: Visual Object and Space Perception battery (Warrington & James, 1991); PALPA: Psycholinguistic Assessment of Language Processing in Aphasia (Kay, et al., 1992b); ⁺ Tests from (Bozeat, Lambon Ralph, et al., 2000).

Table 3. Proportion of different error types for the 10 PA and 10 SD cases. Standard deviations are provided in parentheses.

		HFR*	LFR*	HFE*	LFE*	Total [†]
Omission Errors	PA	0 (0)	0.017 (0.017)	0 (0)	0.038 (0.017)	0.021 (0.034) [‡]
	SD	0 (0)	0 (0)	0 (0)	0 (0)	0 (0) [‡]
LARC Errors	PA	0.017 (0.017)	0.096 (0.052)	0.248 (0.092)	0.268 (0.042)	0.198 (0.111) [‡]
	SD	0.31 (0.098)	0.142 (0.056)	0.36 (0.107)	0.547 (0.054)	0.434 (0.203) [‡]
Visual Errors	PA	0.213 (0.066)	0.365 (0.081)	0.191 (0.051)	0.288 (0.048)	0.279 (0.152) [#]
	SD	0.139 (0.074)	0.193 (0.07)	0.289 (0.081)	0.17 (0.033)	0.187 (0.128) [#]
Letter Omissions	PA	0.052 (0.034)	0.161 (0.057)	0.185 (0.074)	0.097 (0.018)	0.122 (0.067)
	SD	0.028 (0.021)	0.137 (0.069)	0.16 (0.068)	0.074 (0.018)	0.086 (0.065)
Letter Additions	PA	0.035 (0.019)	0.066 (0.035)	0.008 (0.008)	0 (0)	0.021 (0.021)
	SD	0.089 (0.05)	0.006 (0.006)	0.022 (0.015)	0.014 (0.007)	0.029 (0.019)
Letter Transpositions	PA	0.112 (0.054)	0.013 (0.009)	0 (0)	0.004 (0.004)	0.024 (0.022) [‡]
	SD	0.061 (0.036)	0 (0)	0 (0)	0 (0)	0.007 (0.012) [‡]
Letter Substitutions	PA	0.332 (0.098)	0.258 (0.076)	0.34 (0.092)	0.166 (0.044)	0.242 (0.07) [‡]
	SD	0.177 (0.075)	0.446 (0.075)	0.125 (0.04)	0.117 (0.016)	0.175 (0.051) [‡]
Other Errors	PA	0.24 (0.129)	0.024 (0.01)	0.029 (0.015)	0.139 (0.034)	0.093 (0.069)
	SD	0.197 (0.1)	0.077 (0.032)	0.044 (0.021)	0.078 (0.015)	0.083 (0.054)

Note: *Proportions = error type/errors per condition. [†]Proportion = error type/total errors. [‡]Significant group difference at p<.05, one-tailed; [#]Marginally significant group difference at p<.08, one-tailed.

Table 4. Proportion of errors of commission that were phonologically identical to another known word for the 10 PA and 10 SD cases. Standard deviations are provided in parentheses.

	HFR*	LFR*	HFE*	LFE*	Total*
PA	0.823 (0.1)	0.903 (0.04)	0.826 (0.062)	0.649 (0.057)	0.777 (0.084) [†]
SD	0.665 (0.1)	0.556 (0.063)	0.43 (0.077)	0.411 (0.039)	0.46 (0.111) [†]


Note: *Proportions = word errors/commission errors per condition. †Proportion = word errors/total commission errors. ‡Significant group difference at $p < .000005$, one-tailed.

For Peer Review Only


APPENDIX

Structural scans for eight patients in the present study where these were available. Patients are ordered from mildest to most severe ordered from least to most impaired according to high frequency regular word reading accuracy.


Patient JW




Patient JM




Patient TS



Patient SC




Patient JWF




Patient MS



Patient AT



Patient EW



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

REFERENCES

- 1
2
3
4
5
6
7 Acosta-Cabronero, J., Patterson, K., Fryer, T., Hodges, J., Pengas, G., Williams, G., & Nestor, P.
8 (2011). Atrophy, hypometabolism and white matter abnormalities in semantic dementia
9 tell a coherent story. *brain*, 134, 2025-2035.
- 10 Adlam, A. L. R., Patterson, K., Rogers, T. T., Nestor, P. J., Salmond, C. H., Acosta-Cabronero, J., &
11 Hodges, J. R. (2006). Semantic dementia and fluent primary progressive aphasia: Two sides
12 of the same coin? *Brain*, 129(11), 3066-3080.
- 13 Arguin, M., & Bub, D. N. (1993). Single-character processing in a case of pure alexia.
14 *Neuropsychologia*, 31(5), 435-458.
- 15 Arguin, M., Fiset, S., & Bub, D. (2002). Sequential and parallel letter processing in letter-by-letter
16 dyslexia. *Cognitive Neuropsychology*, 19(6), 535-555.
- 17 Bailey, T. M., & Hahn, U. (2005). Phoneme similarity and confusability. *Journal of Memory and*
18 *Language*, 52(3), 347-370.
- 19 Behrmann, M., Nelson, J., & Sekuler, E. B. (1998). Visual complexity in letter-by-letter reading:
20 'Pure' alexia is not pure. *Neuropsychologia*, 36(11), 1115-1132.
- 21 Behrmann, M., & Plaut, D. C. (2013a). Bilateral Hemispheric Processing of Words and Faces:
22 Evidence from Word Impairments in Prosopagnosia and Face Impairments in Pure Alexia.
23 *Cereb Cortex*. doi: 10.1093/cercor/bhs390
- 24 Behrmann, M., & Plaut, D. C. (2013b). Distributed circuits, not circumscribed centers, mediate
25 visual recognition. *Trends Cogn Sci*, 17(5), 210-219. doi: 10.1016/j.tics.2013.03.007
- 26 Behrmann, M., Plaut, D. C., & Nelson, J. (1998). A literature review and new data supporting an
27 interactive account of letter-by-letter reading. *Cognitive Neuropsychology*, 15(1-2), 7-51.
- 28 Behrmann, M., & Shallice, T. (1995). Pure Alexia - a Nonspatial Visual Disorder Affecting Letter
29 Activation. *Cognitive Neuropsychology*, 12, 409-454.
- 30 Blazely, A. M., Coltheart, M., & Casey, B. J. (2005). Semantic impairment with and without surface
31 dyslexia: Implications for models of reading. *Cognitive Neuropsychology*, 22(6), 695-717.
- 32 Bouma, H. (1971). Visual recognition of isolated lower-case letters. *Vision Research*, 11(5), 459-
33 474.
- 34 Bozeat, S., Gregory, C. A., Lambon Ralph, M. A., & Hodges, J. R. (2000). Which neuropsychiatric and
35 behavioural features distinguish frontal and temporal variants of frontotemporal
36 dementia from Alzheimer's disease? *Journal of Neurology Neurosurgery and Psychiatry*,
37 69(2), 178-186.
- 38 Bozeat, S., Lambon Ralph, M. A., Patterson, K., Garrard, P., & Hodges, J. R. (2000). Non-verbal
39 semantic impairment in semantic dementia. *Neuropsychologia*, 38(9), 1207-1215.
- 40 Bub, D. N., Black, S., & Howell, J. (1989). Word recognition and orthographic context effects in a
41 letter- by-letter reader. *Brain and Language*, 36, 357-376.
- 42 Chang, Y. N., Furber, S., & Welbourne, S. (2012a). Modelling normal and impaired letter
43 recognition: Implications for understanding pure alexic reading. *Neuropsychologia*, 50(12),
44 2773-2788.
- 45 Chang, Y. N., Furber, S., & Welbourne, S. (2012b). " Serial" effects in parallel models of reading.
46 *Cognitive Psychology*, 64(4), 267-291.
- 47 Cipelotti, L., & Warrington, E. K. (1995). Semantic memory and reading abilities: a case report.
48 *Journal of the International Neuropsychological Society : JINS*, 1(1), 104-110.
- 49 Cohen, L., Lehericy, S., Chochon, F., Lemer, C., Rivaud, S., & Dehaene, S. (2002). Language-specific
50 tuning of visual cortex? Functional properties of the Visual Word Form Area. *Brain*, 125(5),
51 1054-1069.
- 52 Coltheart, M., Tree, J. J., & Saunders, S. J. (2010). Computational Modeling of Reading in Semantic
53 Dementia: Comment on Woollams, Lambon Ralph, Plaut, and Patterson (2007).
54 *Psychological Review*, 117(1), 256-271.
- 55
56
57
58
59
60


- 1
2
3 Cumming, T. B., Patterson, K., Verfaellie, M. M., & Graham, K. S. (2006). One bird with two stones:
4 Abnormal word length effects in pure alexia and semantic dementia. *Cognitive*
5 *Neuropsychology*, 23(8), 1130-1161.
- 6 Déjerine, J. (1892). Contribution a l'étude anatomo-pathologique et clinique des différentes
7 variétés de cécité-verbale. *Mémoires Société Biologique*, 4, 61-90.
- 8 Farah, M. J., & McClelland, J. L. (1991). A computational model of semantic memory impairment:
9 modality specificity and emergent category specificity. *J Exp Psychol Gen*, 120(4), 339-357.
- 10 Farah, M. J., & Wallace, M. (1991). Pure alexia as a visual impairment: A reconsideration. *Cognitive*
11 *Neuropsychology*, 8(313-334).
- 12 Fiset, D., Arguin, M., Bub, D., Humphreys, G. W., & Riddoch, M. J. (2005). How to make the word-
13 length effect disappear in letter-by-letter dyslexia: Implications for an account of the
14 disorder. *Psychological Science*, 16(7), 535-541.
- 15 Fiset, S., Arguin, M., & Fiset, D. (2006). An attempt to simulate letter-by-letter dyslexia in normal
16 readers. *Brain and Language*, 98(3), 251-263.
- 17 Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-mental state: A practical method for
18 grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*,
19 12(3), 189-198.
- 20 Friedman, R. B., & Alexander, M. P. (1984). Pictures, Images, and Pure Alexia - a Case-Study.
21 *Cognitive Neuropsychology*, 1(1), 9-23. doi: Doi 10.1080/02643298408252014
- 22 Gold, B. T., Balota, D. A., Cortese, M. J., Sergent-Marshall, S. D., Snyder, A. Z., Salat, D. H., . . .
23 Buckner, R. L. (2005). Differing neuropsychological and neuroanatomical correlates of
24 abnormal reading in early-stage semantic dementia and dementia of the Alzheimer type.
25 *Neuropsychologia*, 43(6), 833-846.
- 26 Graham, K. S., Patterson, K., & Hodges, J. R. (1995). Progressive pure anomia: Insufficient
27 activation of phonology by meaning. *Neurocase* 1(25-38).
- 28 Graham, N. L., Patterson, K., & Hodges, J. R. (2000). The impact of semantic memory impairment
29 on spelling: Evidence from semantic dementia. *Neuropsychologia*, 38(2), 143-163.
- 30 Hanley, J. R., & Kay, J. (1996). Reading speed in pure alexia. *Neuropsychologia*, 34, 1165-1174.
- 31 Harm, M. W., & Seidenberg, M. S. (2004). Computing the meanings of words in reading:
32 Cooperative division of labor between visual and phonological processes. *Psychological*
33 *Review*, 111(3), 662-720.
- 34 Harris, L., Olson, A., & Humphreys, G. (2013). Overcoming the effect of letter confusability in
35 letter-by-letter reading: a rehabilitation study. *Neuropsychol Rehabil*, 23(3), 429-462. doi:
36 10.1080/09602011.2013.776500
- 37 Hasson, U., Harel, M., Levy, I., & Malach, R. (2003). Large-scale mirror-symmetry organization of
38 human occipito-temporal object areas. *Neuron*, 37(6), 1027-1041.
- 39 Hasson, U., Levy, I., Behrmann, M., Hendler, T., & Malach, R. (2002). Eccentricity bias as an
40 organizing principle for human high-order object areas. *Neuron*, 34(3), 479-490. doi: Doi
41 10.1016/S0896-6273(02)00662-1
- 42 Henderson, L. (1982). *Orthography and word recognition in reading*. London: Academic Press.
- 43 Hodges, J. R., Patterson, K., Oxbury, S., & Funnell, E. (1992). Semantic dementia. Progressive fluent
44 aphasia with temporal lobe atrophy. *Brain*, 115(6), 1783-1806.
- 45 Hodges, J. R., Salmon, D. P., & Butters, N. (1992). Semantic memory impairment in Alzheimer's
46 disease: Failure of access or degraded knowledge? *Neuropsychologia*, 30(4), 301-314.
- 47 Howard, D. (1991). Letter-by-letter readers: Evidence for parallel processing. In D. Besner & G. W.
48 Humphreys (Eds.), *Basic Processes in Reading: Visual word recognition*. Hove: Lawrence
49 Erlbaum.
- 50 Howard, D., & Patterson, K. (1992). *Pyramids and palmtrees: A test of semantic access from*
51 *pictures and words*. Bury St. Edmunds: Thames Valley Test Company.
- 52
53
54
55
56
57
58
59
60

- Ivnik, R. J., Malec, J. F., Smith, G. E., Tangalos, E. G., Petersen, R. C., Kokmen, E., & Kurland, L. T. (1992). Mayo's Older Americans Normative Studies: WAIS-R norms for ages 56 to 97. *Clinical Neuropsychologist*, 6(SUPPL.), 1-30.
- Jefferies, E., Patterson, K., Jones, R. W., & Lambon Ralph, M. A. (2009). Comprehension of concrete and abstract words in semantic dementia. *Neuropsychology*, 23(4), 492-499. doi: 10.1037/a0015452
- Jefferies, E., Patterson, K., & Lambon Ralph, M. A. (2008). Deficits of knowledge versus executive control in semantic cognition: Insights from cued naming. *Neuropsychologia*, 46(2), 649-658.
- Johnson, R. L., & Rayner, K. (2007). Top-down and bottom-up effects in pure alexia: evidence from eye movements. *Neuropsychologia*, 45(10), 2246-2257. doi: 10.1016/j.neuropsychologia.2007.02.026
- Kay, J., & Hanley, R. (1991). Simultaneous Form Perception and Serial Letter Recognition in a Case of Letter-by-Letter Reading. *Cognitive Neuropsychology*, 8, 249-273.
- Kay, J., Lesser, R., & Coltheart, M. (1992a). Psycholinguistic Assessments of Language Processing in Aphasia (PALPA). *Hove: Erlbaum*.
- Kay, J., Lesser, R., & Coltheart, M. (1992b). Psycholinguistic Assessments of Language Processing in Aphasia (PALPA): An introduction. *Aphasiology*, 10, 159-180.
- Knott, R., Patterson, K., & Hodges, J. R. (1997). Lexical and Semantic Binding Effects in Short-term Memory: Evidence from Semantic Dementia. *Cognitive Neuropsychology*, 14(8), 1165-1216.
- Knott, R., Patterson, K., & Hodges, J. R. (2000). The role of speech production in auditory-verbal short-term memory: Evidence from progressive fluent aphasia. *Neuropsychologia*, 38(2), 125-142.
- Kučera, H., & Francis, W. N. (1967). *Computational analysis of present-day American English*. Providence, RI: Brown University Press.
- Lambon Ralph, M. A. L., Graham, K. S., Ellis, A. W., & Hodges, J. R. (1998). Naming in semantic dementia - What matters? *Neuropsychologia*, 36(8), 775-784.
- Levy, I., Hasson, U., Avidan, G., Hendler, T., & Malach, R. (2001). Center-periphery organization of human object areas. *Nat Neurosci*, 4(5), 533-539. doi: 10.1038/87490
- Lezak, M. (1976). *Neuropsychological Assessment*. New York: Oxford University Press.
- Malach, R., Levy, I., & Hasson, U. (2002). The topography of high-order human object areas. *Trends in Cognitive Sciences*, 6(4), 176-184.
- Marshall, J. C., & Newcombe, F. (1973). Patterns of Paralexia - Psycholinguistic Approach. *Journal of Psycholinguistic Research*, 2, 175-199.
- Mion, M., Patterson, K., Acosta-Cabronero, J., Pengas, G., Izquierdo-Garcia, D., Hong, Y. T., . . . Nestor, P. J. (2010). What the left and right anterior fusiform gyri tell us about semantic memory. *Brain*, 133(11), 3256-3268.
- Miozzo, M., & Caramazza, A. (1998). Varieties of pure alexia: The case of failure to access graphemic representations. *Cognitive Neuropsychology*, 15(1-2), 203-238.
- Mycroft, R. H., Behrmann, M., & Kay, J. (2009). Visuo-perceptual deficits in letter-by-letter reading? *Neuropsychologia*, 47(7), 1733-1744.
- Near, D., Snowden, J. S., Gustafson, L., Passant, U., Stuss, D., Black, S., . . . Benson, D. F. (1998). Frontotemporal lobar degeneration: A consensus on clinical diagnostic criteria. *Neurology*, 51(6), 1546-1554.
- Nestor, A., Behrmann, M., & Plaut, D. C. (2013). The neural basis of visual word form processing: a multivariate investigation. *Cereb Cortex*, 23(7), 1673-1684. doi: 10.1093/cercor/bhs158
- Nestor, P. J., Fryer, T. D., & Hodges, J. R. (2006). Declarative memory impairments in Alzheimer's disease and semantic dementia. *NeuroImage*, 30(3), 1010-1020.
- Noble, K., Glosser, G., & Grossman, M. (2000). Oral reading in Dementia. *Brain and Language*, 74(1), 48-69.

- 1
2
3 Patterson, K., Graham, N., & Hodges, J. R. (1994). The impact of semantic memory loss on
4 phonological representations. *Journal of Cognitive Neuroscience*, 6(1), 57-69.
- 5 Patterson, K., & Hodges, J. R. (1992). Deterioration of word meaning: Implications for reading.
6 *Neuropsychologia*, 30(12), 1025-1040.
- 7 Patterson, K., & Kay, J. (1982). Letter-by-letter reading: psychological descriptions of a neurological
8 syndrome. *Quarterly Journal of Experimental Psychology Section A: Human Experimental*
9 *Psychology*, 34(Pt 3), 411-441.
- 10 Patterson, K., & Lambon Ralph, M. A. (1999). Selective disorders of reading? *Current Opinion in*
11 *Neurobiology*, 9(2), 235-239.
- 12 Patterson, K., & Marcel, A. J. (1992). Phonological ALEXIA or PHONOLOGICAL alexia?
13 Patterson, K., Ralph, M. A. L., Jefferies, E., Woollams, A., Jones, R., Hodges, J. R., & Rogers, T. T.
14 (2006). "Presemantic" cognition in semantic dementia: Six deficits in search of an
15 explanation. *Journal of Cognitive Neuroscience*, 18(2), 169-183.
- 16 Pelli, D. G., Burns, C. W., Farell, B., & Moore-Page, D. C. (2006). Feature detection and letter
17 identification. *Vision Research*, 46(28), 4646-4674. doi:
18 <http://dx.doi.org/10.1016/j.visres.2006.04.023>
- 19 Perri, R., Bartolomeo, P., & Silveri, M. C. (1996). Letter dyslexia in a letter-by-letter reader. *Brain*
20 *and Language*, 53, 390-407.
- 21 Pflugshaupt, T., Suchan, J., Mandler, M. A., Sokolov, A. N., Trauzettel-Klosinski, S., & Karnath, H. O.
22 (2011). Do patients with pure alexia suffer from a specific word form processing deficit?
23 Evidence from 'wrods with trasnpsoed letetrs'. *Neuropsychologia*, 49(5), 1294-1301.
- 24 Plaut, D. C., & Behrmann, M. (2011). Complementary neural representations for faces and words:
25 A computational exploration. *Cognitive Neuropsychology*, 28(3-4), 251-275.
- 26 Plaut, D. C., McClelland, J. L., Seidenberg, M. S., & Patterson, K. (1996). Understanding Normal and
27 Impaired Word Reading: Computational Principles in Quasi-Regular Domains.
28 *Psychological Review*, 103(1), 56-115.
- 29 Plaut, D. C., & Shallice, T. (1993). Perseverative and semantic influences on visual object naming
30 errors in optic aphasia: A connectionist account. *Journal of Cognitive Neuroscience*, 5(1),
31 89-117.
- 32 Price, C. J., & Devlin, J. T. (2003). The myth of the visual word form area. *Neuroimage*, 19(3), 473-
33 481.
- 34 Price, C. J., & Devlin, J. T. (2011). The interactive account of ventral occipitotemporal contributions
35 to reading. *Trends Cogn Sci*, 15(6), 246-253. doi: 10.1016/j.tics.2011.04.001
- 36 Rapcsak, S. Z., & Beeson, P. M. (2004). The role of left posterior inferior temporal cortex in
37 spelling. *Neurology*, 62(12), 2221-2229.
- 38 Reuter-Lorenz, P. A., & Brunn, J. L. (1990). A Prelexical Basis for Letter-by-Letter Reading - a Case-
39 Study. *Cognitive Neuropsychology*, 7, 1-20.
- 40 Roberts, D. J., Lambon Ralph, M. A., & Woollams, A. M. (2010). When does less yield more? The
41 impact of severity upon implicit recognition in pure alexia. *Neuropsychologia*, 48(9), 2437-
42 2446.
- 43 Roberts, D. J., Woollams, A. M., Kim, E., Beeson, P. M., Rapcsak, S. Z., & Lambon Ralph, M. A.
44 (2013). Efficient Visual Object and Word Recognition Relies on High Spatial Frequency
45 Coding in the Left Posterior Fusiform Gyrus: Evidence from a Case-Series of Patients with
46 Ventral Occipito-Temporal Cortex Damage. *Cerebral Cortex*, 23(11), 2568-2580. doi:
47 10.1093/cercor/bhs224
- 48 Rogers, T. T., Lambon Ralph, M. A., Garrard, P., Bozeat, S., McClelland, J. L., Hodges, J. R., &
49 Patterson, K. (2004). Structure and Deterioration of Semantic Memory: A
50 Neuropsychological and Computational Investigation. *Psychological Review*, 111(1), 205-
51 235.
- 52 Rosazza, C., Appollonio, I., Isella, V., & Shallice, T. (2007). Qualitatively different forms of pure
53 alexia. *Cogn Neuropsychol*, 24(4), 393-418. doi: 10.1080/02643290701377877
- 54
55
56
57
58
59
60

- 1
2
3 Schneider, W., Eschman, A., & Zuccolotto, A. (2002). E-Prime user's guide.
- 4 Starrfelt, R., & Behrmann, M. (2011). Number reading in pure alexia--a review. *Neuropsychologia*,
5 49(9), 2283-2298. doi: 10.1016/j.neuropsychologia.2011.04.028
- 6 Starrfelt, R., Habekost, T., & Gerlach, C. (2010). Visual processing in pure alexia: a case study. [Case
7 Reports]. *Cortex*, 46(2), 242-255. doi: 10.1016/j.cortex.2009.03.013
- 8 Starrfelt, R., Habekost, T., & Leff, A. P. (2009). Too little, too late: reduced visual span and speed
9 characterize pure alexia. *Cereb Cortex*, 19(12), 2880-2890. doi: 10.1093/cercor/bhp059
- 10 Starrfelt, R., Nielsen, S., Habekost, T., & Andersen, T. S. (2013). How low can you go: Spatial
11 frequency sensitivity in a patient with pure alexia. *Brain and Language*, 126(2), 188-192.
12 doi: <http://dx.doi.org/10.1016/j.bandl.2013.05.006>
- 13 Tadros, K., Fiset, D., Gosselin, F., & Arguin, M. (2009). A Medium spatial frequency trough causes
14 letter-by-letter dyslexia in normal readers. *Journal of Vision*, 9(8), 822. doi:
15 10.1167/9.8.822
- 16
17 Vinckier, F., Dehaene, S., Jobert, A., Dubus, J. P., Sigman, M., & Cohen, L. (2007). Hierarchical
18 Coding of Letter Strings in the Ventral Stream: Dissecting the Inner Organization of the
19 Visual Word-Form System. *Neuron*, 55(1), 143-156.
- 20 Vogel, A. C., Petersen, S. E., & Schlaggar, B. L. (2012). The left occipitotemporal cortex does not
21 show preferential activity for words. *Cereb Cortex*, 22(12), 2715-2732. doi:
22 10.1093/cercor/bhr295
- 23 Warrington, E. K., & James, M. (1991). *The Visual Object and Space Perception Battery*. Bury St.
24 Edmunds, Suffolk: Thames Valley Test Company.
- 25 Warrington, E. K., & Langdon, D. (1994). Spelling Dyslexia - a Deficit of the Visual Word-Form.
26 *Journal of Neurology Neurosurgery and Psychiatry*, 57, 211-216.
- 27 Warrington, E. K., & Shallice, T. (1979). Semantic access dyslexia. *Brain*, 102, 43-63.
- 28 Warrington, E. K., & Shallice, T. (1980). Word-form dyslexia. *Brain*, 103(1), 99-112.
- 29 Wechsler, D. A. (1987). *Wechsler Memory Scale - Revised*. New York: Psychological Corporation.
- 30 Weekes, B. S. (1997). Differential effects of number of letters on word and nonword naming
31 latency. *Quarterly Journal of Experimental Psychology Section A: Human Experimental*
32 *Psychology*, 50(2), 439-456.
- 33 Welbourne, S. R., Woollams, A. M., Crisp, J., & Lambon Ralph, M. A. (2011). The role of plasticity-
34 related functional reorganization in the explanation of central dyslexias. *Cognitive*
35 *Neuropsychology*, 28(2), 65-108.
- 36 Woodhead, Z. V. J., Wise, R. J. S., Sereno, M., & Leech, R. (2011). Dissociation of sensitivity to
37 spatial frequency in word and face preferential areas of the fusiform gyrus. *Cerebral*
38 *Cortex*, 21(10), 2307-2312.
- 39 Woollams, A. M., Cooper-Pye, E., Hodges, J. R., & Patterson, K. (2008). Anomia: A doubly typical
40 signature of semantic dementia. *Neuropsychologia*, 46(10), 2503-2514.
- 41 Woollams, A. M., Lambon Ralph, M. A., Plaut, D. C., & Patterson, K. (2007). SD-squared: On the
42 association between semantic dementia and surface dyslexia. *Psychological Review*,
43 114(2), 316-339.
- 44 Woollams, A. M., Lambon Ralph, M. A., Plaut, D. C., & Patterson, K. (2010). SD-Squared Revisited:
45 Reply to Coltheart, Tree, and Saunders (2010). *Psychological Review*, 117(1), 273-281.
- 46 Yong, K. X., Warren, J. D., Warrington, E. K., & Crutch, S. J. (2013). Intact reading in patients with
47 profound early visual dysfunction. *Cortex*. doi: 10.1016/j.cortex.2013.01.009
- 48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Reading aloud accuracy for 10 pure alexic and 10 semantic dementia patients according to frequency and regularity. Error bars represent +/- standard error.



HFR= High Frequency Regular; LFR = Low Frequency Regular; HFE = High Frequency Exception; LFE – Low Frequency Exception




Figure 2. Proportion of LARC and Letter (Visual + Transposition + Substitution) errors for the 10 PA and 10 SD patients. Error bars represent +/- standard error.




Figure 3. Proportion of word and nonword errors for the 10 PA and 10 SD patients. Error bars represent +/- standard error.




Figure 4. Visual similarity of letter substitution errors for the 10 PA (left) and 10 SD (right) patients. Values represent proportion of all substitution errors.




Figure 5. Proportion of consonant substitution errors according to number of phonetic features (place, manner, voice, sonorance) shared between presented and reported phonemes for the 10 PA and 10 SD patients.




Figure 6. Schematic representation of the loci of deficits undermining reading in PA (left) versus SD (right) within a triangle model of reading. Filled ovals represent damaged components, grey ovals represent subsequently disrupted processing.