BMJ Global Health

On the importance of primary and community healthcare in relation to global health and environmental threats: lessons from the COVID-19 crisis

Paolo Lauriola , ¹ Piedad Martín-Olmedo, ² Giovanni S. Leonardi , ³ Catherine Bouland , ⁴ Robert Verheij, ⁵ Michel L A Dückers , ⁵ Martie van Tongeren, ⁶ Ferdinando Laghi, ⁷ Peter van den Hazel, ⁸ Ozden Gokdemir , ⁹ Evelyn Segredo, ¹⁰ Ruth A Etzel, ¹¹ Alan Abelsohn, ¹² Fabrizio Bianchi, ¹³ Roberto Romizi, ¹⁴ Giuseppe Miserotti, ¹⁵ Francesco Romizi, ¹⁴ Paolo Bortolotti, ¹⁶ Emanuele Vinci, ¹⁷ Guido Giustetto, ¹⁸ Mariagrazia Santamaria, ¹⁹ Alice Serafini , ²⁰ Samantha Pegoraro, ²¹ Raymond Agius, ⁶ Ariana Zeka , ²²

To cite: Lauriola P, Martín-Olmedo P, Leonardi GS, et al. On the importance of primary and community healthcare in relation to global health and environmental threats: lessons from the COVID-19 crisis. BMJ Global Health 2021;6:e004111. doi:10.1136/ bmigh-2020-004111

Handling editor Seye Abimbola

Received 6 October 2020 Revised 29 December 2020 Accepted 6 January 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by RM.I

For numbered affiliations see end of article.

Correspondence to

Dr Ariana Zeka; Ariana.Zeka@brunel.ac.uk and Dr Paolo Lauriola; paolo.lauriola@gmail.com

ABSTRACT

In the course of the COVID-19 pandemic, it has become clear that primary healthcare systems play a critical role in clinical care, such as patient screening, triage, physical and psychological support and also in promoting good community advice and awareness in coordination with secondary healthcare and preventive care. Because of the role of social and environmental factors in COVID-19 transmission and burden of disease, it is essential to ensure that there is adequate coordination of populationbased health services and public health interventions. The COVID-19 pandemic has shown the primary and community healthcare (P&CHC) system's weaknesses worldwide. In many instances, P&CHC played only a minor role, the emphasis being on hospital and intensive care beds. This was compounded by political failures, in supporting local community resilience. Placing community building, social cohesion and resilience at the forefront of dealing with the COVID-19 crisis can help align solutions that provide a vision of 'planetary health'. This can be achieved by involving local well-being and participation in the face of any pervasive health and environmental crisis, including other epidemics and large-scale ecological crises. This paper proposes that P&CHC should take on two critical roles: first, to support local problem-solving efforts and to serve as a partner in innovative approaches to safeguarding community well-being; and second, to understand the local environment and health risks in the context of the global health perspective. We see this as an opportunity of immediate value and broad consequence beyond the control of the COVID-19 pandemic.

INTRODUCTION AND OBJECTIVES

The novel SARS-CoV-2 and its resultant COVID-19 disease is the most challenging

Summary box

- ▶ In 1978, the Alma-Ata (Kazakhstan) Declaration asserted the strategic importance of high-quality primary healthcare in the creation of effective and responsive healthcare systems; unfortunately, 42 years later, this vision has not been attained yet.
- ▶ In October 2018, health experts and policymakers met in Astana (Kazakhstan) to renew the commitment for comprehensive available healthcare to all, reaffirming the commitments to the Alma-Ata core principles.
- ► The lack of involvement of the primary and community healthcare (P&CHC) systems during the COVID-19 pandemic has highlighted their crucial role in clinical and public health functions such as diagnosis (screening), triage, contact tracing and in the short-term and long-term physical and psychological monitoring and management of patients.
- ► The current emergency has also shown the need for P&CHC to work in coordination with other healthcare and public health organisations at the community level
- There is an urgent need for environmental public health policies with an intersectoral and global perspective acknowledging the influence of the environment and the need to act at the community level.
- ▶ Besides the traditional focus on the individual, it is essential that public health agencies encourage and promote the involvement of P&CHC within the framework of community and environmental health.
- ► The value of this has been highlighted by the response to COVID-19, but it is equally central to the management of other crises affecting communities.



BMJ Glob Health: first published as 10.1136/bmjgh-2020-004111 on 10 March 2021. Downloaded from http://gh.bmj.com/ on March 12, 2021 at Brunel Uni Consortia. Protected by copyright

health emergency faced by humanity in living memory. This pandemic represents a grave threat to global health, economy and the social well-being of people.

The COVID-19 pandemic has emphasised the crucial role that primary and community healthcare (P&CHC) systems¹² could play in diagnosis or screening and triage, as well as in the short-term and long-term monitoring of the physical and psychological conditions of those affected. The current emergency has also highlighted the need for P&CHC to work on tracing and management of contacts in coordination with other healthcare organisations such as local and national public health institutions, ^{3 4} as well as those at the community level⁵ such as community-based civil society groups and community health workers.

Because of the role of social and environmental factors in COVID-19 transmission and in the burden of disease, it is essential to ensure that population-based health services and interventions are adequately prioritised and coordinated between P&CHC providers and public health organisations.

There are recognisable dimensions to the complex domain of how decisions are made to manage rapidly emerging problems. Each approach to crisis management has its merits, and they are all interconnected. The role of P&CHC is central to the effective management of any health crisis.

Despite historical and ongoing discrepancies between and within the policy and scientific communities, mainly due to the complexity mentioned above, the pandemic has shown the importance of facilitating the interrelationship and mutual support between the health, social, environmental and political-economic domains.

The goal of this article is to highlight the crucial role that P&CHC could play in managing the COVID-19 pandemic, in addressing key environmental and social aspects involved in viral transmission, and assist in the design of the interventions to control it. The article discusses how the success of such interventions depends on an overall vision of integrating environmental health with social and healthcare. It makes the case that enhancing such capacity now will have a more comprehensive benefit for better addressing this crisis and others

We lay emphasis on the importance of a multidisciplinary and effective workforce in environmental public health (EPH), in collaboration with P&CHC and hospitals.

The COVID-19 pandemic is not the only concern we address with this proposed strategy; this is an exemplar of a 'stress test' of global health and environmental threats.

SARS-COV-2 AND COVID-19: AN EXAMPLE OF **ENVIRONMENTAL AND SOCIOPOLITICAL IMBALANCE**

The origin of SARS-CoV-2 and the subsequent COVID-19 pandemic have revealed the precariousness of the systems

on which trade, food, energy, transportation and social safety and security nets depend.

The reported origin of the current pandemic emphasises the exceptional relationship that exists between the environment, animal and human health (One Health approach).8 Evidence suggests that SARS-CoV-2 emerged from transportation and trading of wild animals in wet markets. About 1 billion cases of illness and millions of deaths occur every year from zoonoses, which comprise about 60% of all emerging infectious diseases globally. 10 The majority of these (60.3%) are caused by zoonotic pathogens, out of which 72% have a wildlife origin. 11 Ecosystem health, wildlife health and human health are increasingly interconnected, leading to an urgent need to develop locally and globally coordinated plans for better addressing the impacts of environmental change on zoonotic and parasitic infections, ¹² including COVID-19.¹³

Local, national, large-scale and systemic environmental changes and disruptions, such as biodiversity loss, disruption of natural habitats, extinction of species, climate crisis, pollution, together with societal and demographic changes (eg, urbanisation, increase in mobility, population ageing), are likely determinants of different epidemics.¹⁴ Both environmental and social factors play an important role by facilitating the occurrence of new or re-emerging infectious diseases, and/or by greatly modulating their health impacts.

GLOBAL/LOCAL, TWO SIDES OF THE SAME COIN

The way in which the COVID-19 crisis developed in time and space made clear that the distinction between local and global has become largely unimportant.¹⁵ The local ecological wildlife and human interactions in Wuhan, China, have been recognised as the origin of the SARS-CoV-2 spillover, which afterwards quickly spread across the globe. Such local incidents are related to global environmental issues such as ecological erosion and the climate crisis. Regional and local political and economic decisions that promote unsustainable practices lead to deforestation, intensive farming, unsustainable energy production, air pollution and deterioration of water and soil, all of which contribute further to the global climate crisis, increasing the likelihood of emerging global impacts in the future. The role of a globalised economic order that prioritises short-term profits of private interests, which benefit actors in wealthy countries and elites in low-and-middle-income countries (LMICs), while dismissing environmental and social fallout as 'negative externalities', are clearly recognised. There are many historical examples of how societies did not recognise the 'circularity in nature' with disastrous results. Such lessons have shown how big businesses affect the environment, and what individuals and communities can do about our environmental problems today.¹⁶

And while global system-level change is needed, communities can also play an important role in helping



to address environmental problems at multiple levels. P&CHC can contribute to addressing such global challenges, with the view of prevention and protection from harm of communities, with global benefits by sharing of experiences and expertise to improve competencies and capabilities at the local level.

These efforts are not at all simple to deploy for many reasons such as the overall organisation or conflicts between science and politics, with some clear examples across the world as provided here ^{17–19}

The uneven distribution of COVID-19 cases and severity of the disease²⁰ are determined by socioeconomic circumstances, inequality associated with poor working and living conditions, access to services, mobility and pre-existing health conditions associated with deprivation.²¹ Populations lacking access to health services in normal circumstances are left most vulnerable during times of crisis,²² and addressing such social determinants and inequities is part of the challenge of P&CHC at global level and first and foremost at local level.

P&CHC AS A CORNERSTONE TO ADDRESS GLOBAL ISSUES AT THE LOCAL LEVEL

P&CHC has historically had a crucial role in communities dealing with global challenges. Societies are ageing, with many facing increasing rates of obesity, noncommunicable diseases (NCDs) and new and re-emerging infectious diseases, with health spending rising.²³ It is therefore imperative to redesign health systems that contribute to community health by addressing environmental factors. 16 The rapid spread of COVID-19 added complexity to these challenges. A multidisciplinary P&CHC workforce can play a vital role in recognition and management of environmental and social factors of community health. Family doctors (FDs) and family paediatricians (FPs) are usually the first contact point for patients, even in the current pandemic, especially in high-income countries (see the Low and Middle-Income Countries section for considerations more specific to LMICs).

Health impacts in different settings of P&CHC approach

The main critical threat of the COVID-19 pandemic was likely due to the potential collapse of the health service. This has been extensively investigated worldwide. ^{24 25} The Italian experience is an example.

Italy was the first European country involved in the pandemic and has been one of the worst affected. The first confirmed cases were detected on 31 January 2020. Later, a cluster of cases with the first deaths was identified in -the northern region of Lombardy. The virus had also spread to the neighbouring region of Veneto, but not initially to other areas of the country.

The Italian healthcare system is highly decentralised; different regions implemented different policy responses to COVID-19. The most notable example is the contrast between the approaches adopted by Lombardy and

Veneto, two neighbouring regions with similar socioeconomic profiles.

Lombardy, one of Europe's wealthiest and most industrial productive areas, was disproportionately hit by COVID-19. Veneto, which is similarly wealthy, by contrast, fared significantly better. In the second wave of the pandemic in autumn, Lombardy scored at the highest level of risk. ²⁶ By the end of December 2020, the situation was changing, arguably linked to many contributing factors (virological, organisational), which are under investigation.

The spread of COVID-19 and the policy measures taken in these two regions were shaped by a multitude of factors, including the different public health management choices applied in each region.²⁷ Interventions in Lombardy—like in other regions which experienced much larger case fatality rates—were based mainly on hospital activities.²⁸ The region of Veneto, at least in the first half of the year 2020, relied more on the involvement of primary care doctors, with resulting lower health impacts. While the situation is more complex with many factors (such as socioeconomic, organisational, infrastructure, public health surveillance) contributing to the observed differences in health impacts, this experience highlighted the importance of P&CHC as the front line in dealing with health crises.

Connections of P&CHC

The connection between P&CHC and occupational health (OH) is well established. This is important for better prevention of chronic conditions (such as musculoskeletal or mental health disorders) that lead to absenteeism or early departure from the labour force. Efforts to improve workers' physical and mental health include raising awareness among managers, improving the physical working environment, humanising social relations at work and offering programmes dedicated to encouraging disabled people to return to work. Equally important is the health surveillance of workers especially for physical, chemical or biological hazards at work.

In this respect, COVID-19 presents an almost unique instance where occupational and environmental risk factors have overlapped, generating a health, social and economic impact of a magnitude not seen before. Failure of OH control at work will affect many individuals; failure of control in the community will result in many sick workers. Hence, the close coordination between P&CHC, OH, EPH and enforcement authorities is essential.³⁰

P&CHC can play two critical roles: one, as partners to public health policymakers and agencies in contributing to monitoring impacts and serving as 'anchor institutions' for implementing local solutions and innovative approaches to safeguarding community health; and two, an essential role in community building, social cohesion and resilience. ³²



P&CHC: THE LINCHPIN CONNECTING ENVIRONMENT AND HEALTH

COVID-19 severity and fatality have been shown highest among those made vulnerable by less favourable structural social conditions.²² The unequal distribution of health and mortality burden disfavouring those most vulnerable, between and within countries and communities, shows similarities across the spectrum of health conditions—these being communicable or noncommunicable.^{33 34}

In the case of COVID-19, the term 'syndemic' has been cited, as the health burden has been beyond the number of cases of infection, and has extended to several NCDs aspects. First, biological and social interactions of COVID-19 with underlying noncommunicable health conditions often lead to more serious disease progression, including death. Second, services for NCDs have been considerably disrupted in three-quarters of countries globally with associated impacts. As environmental factors are estimated to account for at least 25% of the preventable burden of disease, including NCDs, a perspective for primary prevention of community disease burden cannot ignore EPH.

The health effects associated with environmental risk factors, usually quantified through mortality, emergency visits and hospitalisations, are underrepresented, as many illnesses treated by FDs or FPs are missed in these numbers.³⁸ For example, vectorborne infectious diseases show a geographical heterogeneity linked to climate crisis.³⁹ In these cases, global strategies addressing climate crisis can and should be implemented locally. The role of local communitybased FDs and FPs is essential in what they can do to encourage mitigation of climate crisis and to assist communities in their adaptation efforts. A clear and comprehensive example of the activities which could be carried out within the P&CHC framework to address global environmental risk factors is provided in box 1.40

IMPLEMENTING STRONGER P&CHC

New approaches can be implemented as preventive health/care measures with a broad integrated perspective. Action is needed at multiple levels, and which is attuned to different contexts.

Local-national level

A policy survey conducted by the Organisation for Economic Co-operation and Development (OECD)¹ suggests that key policies and strategies delivering better P&CHC should be based on:

► New models of care: rather than the single-practice physician, multidisciplinary health teams supported by digital technology should be proactively engaged in preventive care. 41

Box 1 Primary and community healthcare role in relation to global health threats

- ► Public education and raising awareness.
- Day-by-day involvement in local/regional/national strategies to tackle antimicrobial resistance within a One Health approach.
- ► Early alert systems: impending weather extremes, infectious disease outbreaks.
- ▶ Disaster preparedness, including increasing the health system's 'surge' capacity to respond to emergencies.
- Enhanced infectious disease control programmes.
 Food safety, vaccine programmes, case detection and treatment.
- Improved surveillance.

Vector control.

Risk indicators (eg, aeroallergen concentration).

Health outcomes (eg, infectious disease outbreaks, rural suicides, seasonal asthma peaks).

Appropriate health workforce training, including continuing professional development (eg, updated understanding of climatic influences on health, training in public health).

(Adapted from Blashki et al [40]).

- More economic incentives: to encourage P&CHC to work in teams and focus on prevention and continuity of care.
- ▶ A broader role for patients, including involving the patients in the co-management of their health.

Digitalisation is essential to integrate P&CHC further at the community level. 'Precision public health' is an approach which uses data from traditional and emerging sources, for example, big data to target interventions for populations by person, place and time. ⁴² With such approach, supported by strong empirical foundations, electronic health records from P&CHC, especially if in combination with data on environmental circumstances, represent an essential source of data and understanding at the local level. ⁴³

As such, FDs and FPs could play a crucial role in connecting global concerns with local actions through the information they provide, which can influence evidence-based needed policy, in addition to their important role in community awareness and attitudes.⁴⁴

EPH at the level of P&CHC could be strengthened through agreements with:

- 1. Employers of primary care staff (usually in the public sector), to modify contracts and job descriptions of selected currently employed P&CHC staff to identify tasks related to EPH aspects of community health.
- 2. Universities and other agencies required for provision of training of P&CHC staff, to identify appropriate competencies and standards required, and develop related training programmes to enable them to perform the tasks related to EPH aspects of community health. It is possible that a first cohort of P&CHC staff may not have acquired sufficient training in EPH in the absence of formally recognised training programmes, so this aspect will concern successive cohorts; further-

more, lifelong learning would drive towards a larger number.

- 3. Professional registration bodies, such as Ordine dei Medici/General Medical Council and other professional registration authorities, depending on country and type of profession, 45 to conduct and conclude negotiations with employers and universities/training agencies for skills and competencies required to achieve registration of staff who have completed training related to EPH aspects of community health, as part of their professional qualification.
- 4. Health agencies to promote secondary use of electronic health records data from primary and community care facilities and linking that to other data relating to the environment.

International level

International organisations should provide a framework where the crucial role of P&CHC in EPH is emphasised especially with respect to data sharing, responsibilities, capability and consistent capacity building. The following organisations are already dealing with such issues, and should be encouraged to develop such framework with a collaborative perspective.

The World Health Organisation stated that a vision for primary health care (PHC) in the 21st century must be based on individuals and communities as a unique

The World Organization of Family Doctors is an international organisation that aims at improving the quality of life by fostering high standards of care in general practice/family medicine considering the environment.⁴⁷

Health and Environment Alliance is the leading European not-for-profit organisation addressing how the natural and built environments affect health in Europe. 48

International Society of Doctors for the Environment was founded 30 years ago, aiming at educating physicians and the general public.⁴⁹

Low and middle-income countries

Particular attention must be paid to LMICs with populations and economies more vulnerable to the immediate effects of the climate crisis, including zoonotic infections or parasitic disease, and the long-term health, social and economic impacts of the current pandemic, and more generally to the difficulties towards a sustainable and equitable development.

Integration of PHC in view of a community health perspective in LMICs is complex. As such, two points¹ should be raised:

1. Data: many LMICs lack sufficient data on PHC performance, with some countries lacking even basic vital registration data.⁵⁰

Data gaps that influence PHC systems can be found at any level and dimension of the health system, including aspects of governance, leadership and population health management.

2. Integration: integrating a PHC system is a combined effort, and there is no single tool to achieve this. The PHC systems design approach must be strengthened, incorporating human resources for health, financing, facilities, private sector and demand to develop and enhance integrated systems.

P&CHC is mainly based on the awareness of the socioeconomic, capacity and cultural/historical features in which the FDs work. As such, according to the Astana Declaration, ⁵¹ a holistic approach must integrate efforts to deal with other primary conditions such as malnutrition, waterborne diseases and infectious diseases which disproportionately impact LMICs.

According to OECD, in LMICs, attention needs to be directed to:

- District management systems with consistent investments in front-line PHC and community health workers.
- Mobilisation of sufficient financial resources to reduce inequalities in the ability to pay P&CHC services, and to provide financial protection against impoverishment from catastrophic healthcare costs.
- Improved learning between healthcare systems and integrated care, and development of strategies for engaging the private sector.

CAPABILITY AND CAPACITY: THE ROLE OF EDUCATION Background

Knowledge and recognition of EPH issues are not common among physicians and other health practitioners. Physicians, in particular FDs and FPs, should engage with environmental concerns in both their role as clinicians as well as their role as public health practitioners. EPH training of physicians should include knowledge and skills in recognising, diagnosing and treating health problems caused by environmental risk factors. Training would include toxicology and epidemiology, both as basic science, and in the clinical training of FDs before, during and after their specialist training. Education and training of physicians as public health practitioners must include knowledge and skills in general public health, EPH and citizen participation. The successful response of the health system in the Republic of Korea has been attributed to the holistic education of the physicians of that country.²⁵

Proposals and perspectives

The challenge of creating a curriculum that fits a diversity of specialties and levels of training, and covers both the science and implementation of EPH, is not straightforward. However, it is important to be strongly focused on pursuing exactly that.⁵² The overall organisation of this curriculum can be based on an accredited system, according to the different levels and educational needs. It will need to consider flexibly different models of programmes, training and capacity provisions. It will need to be promoted at the highest level of medical

BMJ Glob Health: first published as 10.1136/bmjgh-2020-004111 on 10 March 2021. Downloaded from http://gh.bmj.com/ on March 12, 2021 at Brunel Uni Consortia. Protected by copyright

schools and universities, as well as in community health training (see the Local-National Level section). Universities will need to collaborate with municipal and regional institutions for community health, in identifying standards for required training, at undergraduate and postgraduate levels, in both clinical and community or public health aspects of environmental health. The integration of academic and practical experience is crucial in the success of training in the environmental aspects of clinical medicine and of EPH at the community level. Education must be inclusive, flexible, ongoing and reflective of changes in societal needs.

COVID-19 illustrates very well how the principle of capacity building founded on modification of tasks for P&CHC staff already in post, as we advocate in the Local-National Level section, may be relevant to deployment of existing staff to work on any public health topic. During the COVID-19 emergency, education of all healthcare staff was effective because most of them were already in post. Staff redeployment to COVID-19 duties was usually accepted without modification of contracts and job descriptions, in view of the acceptance of an obvious emergency. A brief statement requiring tasks to be provided on public health, and in particular EPH, should be included in a contract and job description to support and facilitate effective delivery of such tasks by P&CHC staff.

CONCLUSIONS AND RECOMMENDATIONS

There are multiple connections between the environment and health. There is an urgent need for EPH policies with an intersectoral and global perspective.

There is a close relationship between planetary fragility, social frailty and poverty; social justice and environmental justice are two sides of the same coin.

The crucial challenge is to move towards a system that is economically viable, ecologically sustainable and socially fair.⁵³

In the past, the P&CHC was based on the relationship between doctor and patient. Today, the community dimension of the COVID-19 crisis with its NCDs and social aspects provides a lesson for any pervasive crisis.³³ The influence of the environment on health and well-being in their acute and chronic manifestations, and the need to act at the community level are increasingly evident.

Because they are embedded in and closely connected to local communities, primary and community care needs to be strengthened to play a more important role in the relation between individual health and the environment. Global issues need to tangibly translate to local awareness in determining local healthcare policies as well as pandemic and other responses, including the provision of essential data and information about local health. More than anyone else in the healthcare system, P&CHC services are the ones that can make the link between people's health and the environment they live in.

Author affiliations

¹International Society of Doctors for the Environment (ISDE-Italy), Modena, Italy ²Escuela Andaluza de Salud Pública, Granada, Spain

³Department of Public Health, Environments and Society, London School of Hygiene and Tropical Medicine, London, UK

⁴ULB-Ecole de Santé Publique, Bruxelles, Belgium

⁵Netherlands Institute for Health Services Research (NIVEL), Utrecht, The Netherlands

⁶Centre for Occupational and Environmental Health, University of Manchester, Manchester, UK

⁷International Society of Doctors for the Environment (ISDE-Int'l), Geneva. Switzerland

8International Network on Children's Health, Environment and Safety (INCHES), Ellecom, The Netherlands

⁹Faculty of Medicine, Izmir University of Economics, Balcova/Izmir, Turkey

¹⁰Uruguayan Society of Family and Community Medicine, Montevideo, Uruguay

¹¹Milken Institute School of Public Health, The George Washington University, Washington, DC, USA

¹²Department of Family and Community Medicine, University of Toronto, Toronto, Ontario, Canada

¹³Istituto Fisiologia Clinica, Consiglio Nazionale della Ricerca (CNR-IFC), Pisa, Italy

¹⁴International Society of Doctors for the Environment (ISDE-Italy), Arezzo, Italy

¹⁵International Society of Doctors for the Environment (ISDE-Italy), Piacenza, Italy ¹⁶Medical Order Trento, Trento, Italy

¹⁷Health and Environment Working Group, National Medical Orders Federation, Rome, Italy

18 National Medical Orders Federation, Rome, Italy

¹⁹Local Health Authority, Foggia, Italy

²⁰Local Health Authority, Modena, Italy

²¹Italian Climate Network, Milano, Italy

²²Institute of Environment, Health and Societies, Brunel University London, Uxbridge, UK

Twitter Ozden Gokdemir @OzdenGokdemir

Acknowledgements Authors wish to thank the reviewers for their constructive comments which helped improve this paper.

Contributors Coordination: PL. GL. AZ. Conceptualisation of ideas: all authors. Original draft preparation: all authors.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement There are no data in this work.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

Paolo Lauriola http://orcid.org/0000-0003-4768-6612 Giovanni S. Leonardi http://orcid.org/0000-0001-7477-1762 Catherine Bouland http://orcid.org/0000-0003-3085-0543 Michel L A Dückers http://orcid.org/0000-0001-7746-053X Ozden Gokdemir http://orcid.org/0000-0002-0542-5767 Alice Serafini http://orcid.org/0000-0002-7396-2839 Ariana Zeka http://orcid.org/0000-0002-9570-8831

REFERENCES

- OECD Realising the Potential of PHC. OECD health policy studies. Paris: OECD Publishing, 2020.
- Li DKT. Challenges and responsibilities of family doctors in the new global coronavirus outbreak. Fam Med Community Health 2020;8:e000333.



- 3 Li X, Krumholz HM, Yip W, Yip M, et al. Quality of primary health care in China: challenges and recommendations. Lancet 2020:395:1802–12
- 4 Harding-Edgar L, McCartney M, Pollock AM. Test and trace strategy has overlooked importance of clinical input, clinical oversight and integration. J R Soc Med 2020;113:428–32.
- 5 Goodman RA, Bunnell R, Posner SF. What is "community health"? Examining the meaning of an evolving field in public health. *Prev Med* 2014;67 Suppl 1:S58–61.
- 6 Sturmberg JP, Martin CM. COVID-19 how a pandemic reveals that everything is connected to everything else. J Eval Clin Pract 2020;26:1361–7.
- 7 Kreis IA, Leonardi GS, Murray VSG. PHEEDUNET document 1 (16 Pages). Definitions. A European network for the training and development of public health (environment) physicians. EU Public Health Executive Agency project NO: 2006335, 2009. Available: https://www.researchgate.net/publication/336217074_PHEEDUNET_Document_1_Definitions_16_pages_A_EUROPEAN_NETWORK_FOR_THE_TRAINING_AND_DEVELOPMENT_OF_PUBLIC_HEALTH_ENVIRONMENT_PHYSICIANS
- 8 Rabinowitz PM, Pappaioanou M, Bardosh KL, et al. A planetary vision for one health. BMJ Glob Health 2018;3:e001137.
- 9 Aguirre AA, Catherina R, Frye H, et al. Illicit wildlife trade, wet markets, and COVID-19: preventing future pandemics. World Medical & Health Policy 2020;12:256–65.
- 10 Karesh WB, Dobson A, Lloyd-Smith JO, et al. Ecology of zoonoses: natural and unnatural histories. Lancet 2012;380:1936–45.
- 11 Jones KE, Patel NG, Levy MA, et al. Global trends in emerging infectious diseases. *Nature* 2008;451:990–3.
- Hueffer K, Parkinson AJ, Gerlach R, et al. Zoonotic infections in Alaska: disease prevalence, potential impact of climate change and recommended actions for earlier disease detection, research, prevention and control. Int J Circumpolar Health 2013;72. doi:10.3402/ijch.v72i0.19562. [Epub ahead of print: 07 02 2013].
- 13 O'Reilly KM, Auzenbergs M, Jafari Y, et al. Effective transmission across the globe: the role of climate in COVID-19 mitigation strategies. Lancet Planet Health 2020;4:e172.
- 14 McMichael AJ. Globalization, climate change, and human health. N Engl J Med 2013;368:1335–43.
- 15 Shaw R, Kim Y-kyun, Hua J. Governance, technology and citizen behavior in pandemic: lessons from COVID-19 in East Asia. *Progress in Disaster Science* 2020;6:100090.
- 16 Watts N, Amann M, Arnell N, et al. The 2020 report of the Lancet countdown on health and climate change: responding to converging crises. Lancet 2021;397:1–42.
- 17 The Lancet. COVID-19 in Brazil: "So what?". Lancet 2020;395:1461.
- 18 Dyer O. Trump claims public health warnings on covid-19 are a conspiracy against him. *BMJ* 2020;368: :m941.
- 19 Türk Tabipleri Birliği. COVID-19 İzleme Kurulu COVID-19 Pandemisi 4. Ay Değerlendirme Raporu. Ankara 2020. Available: https://www. ttb.org.tr/kutuphane/covid19-rapor_4.pdf
- 20 Williamson EJ, Walker AJ, Bhaskaran K, et al. "OpenSAFELY: factors associated with COVID-19 death in 17 million patients. Nature;584:430–6.
- 21 Gehanno JF, Bonneterre V, Andujar P, et al. How should data on airborne transmission of SARS-CoV-2 change occupational health guidelines? Occup Environ Med 2020;77:736.
- 22 Pirisi A. Low health literacy prevents equal access to care. Lancet 2000;356:356.
- 23 OECD/European Union. Health at a glance: Europe 2020: state of health in the EU cycle. Paris: OECD Publishing, 2020.
- 24 OECD. OECD beyond containment: health systems responses to COVID-19 in the OECD, 2020. Available: http://www.oecd. org/coronavirus/policy-responses/beyond-containment-healthsystems-responses-to-covid-19-in-the-oecd-6ab740c0/
- 25 Palaniappan A, Dave U, Gosine B. Comparing South Korea and Italy's healthcare systems and initiatives to combat COVID-19. Rev Panam Salud Publica 2020;44:e53:1.
- 26 IL Ministro Della Salute. Ordinanza Ministro DELLA salute, 2020. Available: http://www.salute.gov.it/imgs/C_17_notizie_5171_0_file.pdf
- 27 Murgia V, Romizi F, Romizi R. Lauriola, family doctors, environment and COVID-19 in Italy: experiences, suggestions and proposals. BMJ 2020;368:m627.
- 28 Ferrara P, Albano L. COVID-19 and healthcare systems: what should we do next? *Public Health* 2020;185:1–2.
- 29 James C, Devaux M, Sassi F. "Inclusive growth and health", OECD health working papers, no. 103. Paris: OECD Publishing, 2017.
- 30 Agius RM, Robertson JFR, Kendrick D, et al. Covid-19 in the workplace. *BMJ* 2020;370:m3577.

- 31 Koh HK, Bantham A, Geller AC, et al. Anchor institutions: best practices to address social needs and social determinants of health. Am J Public Health 2020;110:309–16.
- 32 Castleden M, McKee M, Murray V, et al. Resilience thinking in health protection. J Public Health 2011;33:369–77.
- 33 McNamara CL, Balaj M, Thomson KH, et al. The socioeconomic distribution of non-communicable diseases in Europe: findings from the European social survey (2014) special module on the social determinants of health. Eur J Public Health 2017;27:22–6.
- 34 Liang D, Shi L, Zhao J, et al. Urban air pollution may enhance COVID-19 case-fatality and mortality rates in the United States. medRxiv.
- 35 Horton R. Offline: COVID-19 is not a pandemic. *The Lancet* 2020;874.
- 36 WHO. The impact of the COVID-19 pandemic on noncommunicable disease resources and services: results of a rapid assessment. Geneva: World Health Organization, 2020.
- 37 Prüss-Üstün A, Jennyfer Wolf J, Corvalán C, et al. Preventing disease through healthy environments: a global assessment of the burden of disease from environmental risks. World Health organization, 2016. Available: https://apps.who.int/iris/handle/10665/ 204585
- 38 Lauriola P, Pegoraro S, Serafini A. The role of general practices for monitoring and protecting the environment and health. results and proposals of the italian project aimed at creating an "italian network of sentinel physicians for the environment" (rimsa) within an international perspective. J Family Med Community Health 2018;5:1160.
- 39 Schaffner F, Mathis A. Dengue and dengue vectors in the who European region: past, present, and scenarios for the future. *Lancet Infect Dis* 2014;14:1271–80.
- 40 Blashki G, Abelsohn A, Woollard R, et al. General Practitioners' responses to global climate change - lessons from clinical experience and the clinical method. Asia Pac Fam Med 2012;11:6.
- 41 Murray CJL, Alamro NMS, Hwang H, et al. Digital public health and COVID-19. Lancet Public Health 2020;5:e469–70.
- 42 Rasmussen SA, Khoury MJ, Del Rio C. Precision public health as a key tool in the COVID-19 response. *JAMA* 2020;324:933–4.
- 43 Schweikardt C, Verheij RA, Donker GA, et al. The historical development of the Dutch sentinel general practice network from a paper-based into a digital primary care monitoring system. J Public Health 2016;24:545–62.
- 44 Xie E, de Barros EF, Abelsohn A, et al. Challenges and opportunities in planetary health for primary care providers. The Lancet Planetary Health 2018;2:e185–7.
- 45 Kreis IA, Murray V, Leonardi G, et al. PHEEDUNET document 4 (47 Pages): curricula for registration (procedures for registration as specialist in environmental public health). European Union public health executive agency project NO: 2006335, 2010. https://www. researchgate.net/publication/336217728_PHEEDUNET_Document_ 4_47_pages_Registration_procedures_for_registration_as_specialist_ in_environmental_public_health_European_Union_Public_Health_ Executive_Agency
- 46 WHO. A vision for PHC in the 21st century: towards universal health coverage and the sustainable development goals. Geneva: World Health Organization and the United Nations Children's Fund (UNICEF), 2018.
- 47 WONCA. WONCA Working Party: environment, 2014. Available: https://www.globalfamilydoctor.com/groups/WorkingParties/ Environment.aspx
- 48 Health and environment alliance (HEAL), 2021. Available: https:// www.env-health.org/
- 49 Doctors for the environment. Doctors for the environment, 2017. Available: http://www.isde.org/
- 50 Veillard J, Cowling K, Bitton A, et al. Better measurement for performance improvement in low- and middle-income countries: the primary health care performance initiative (PHCPI) experience of conceptual framework development and indicator selection. Milbank Q 2017;95:836–83.
- 51 WHO. Global conference on primary health care from Alma-Ata towards universal health coverage and the sustainable development goals, Astana, Kazakhstan 25 and 26, 2018. Available: https://www. who.int/docs/default-source/primary-health/declaration/gcphcdeclaration.pdf
- 52 Omrani OE, Dafallah A, Paniello Castillo B, et al. Envisioning planetary health in every medical curriculum: an international medical student organization's perspective. Med Teach 2020;42:1107–11.
- 53 WHO. Manifesto for a healthy recovery from COVID-19, 2020. Available: https://reliefweb.int/report/world/who-manifesto-healthy-recovery-covid-19